

Montana Natural Heritage - SOC Report

Plant Species of Concern

Species List Last Updated **02/19/2021**

453 Species of Concern

90 Potential Species of Concern

All Records (no filtering)



A program of the Montana State Library's
Natural Resource Information System
operated by the University of Montana.

Introduction

The Montana Natural Heritage Program (MTNHP) serves as the state's information source for Species of Concern (SOC) -- plants and animals that are rare, threatened, and/or have declining populations and as a result are at risk or potentially at risk of extirpation in Montana. This report is based on information gathered from field inventories, publications, reports, herbaria specimens, and the knowledge of botanists and other taxonomic experts. Taxa in the SOC category generally include all vascular plant taxa ranked S1, S2, S3 or SH. Nonvascular taxa (bryophytes and lichens) which are not as well documented or studied as vascular plant taxa in the state, are listed as SOC using similar criteria as vascular taxa but are more strictly limited to those taxa which are believed to be the rarest or most vulnerable to extirpation based on current information.

Designation as a Species of Concern is not a statutory or regulatory classification. Instead, these designations provide a basis for resource managers and decision-makers to make proactive decisions regarding species conservation and data collection priorities in order to maintain viable populations and avoid extirpation of species from the state. MTNHP may designate additional taxa as Potential Species of Concern (PSOC). Taxa in this designation include species or subspecies which may be rare, have a restricted range in the state or are otherwise vulnerable to extirpation in at least part of their range but otherwise do not meet the criteria for inclusion as a SOC. An additional designation of Status Under Review is used for those taxa for which additional information is needed to accurately assign a status rank or for which conflicting information exists. Taxa designated as Status Under Review are not included in this document but can be found in the on-line **Fieldguide** (fieldguide.mt.gov).

This web-based report, which replaces the 2006 Plant Species of Concern publication, identifies vascular plant Species of Concern (SOC), bryophyte SOC and lichen SOC in Montana. The MTNHP continuously reviews and updates status ranks as new information and data become available through field surveys, research, and submitted observations. Status ranks and information supporting them are reviewed by botanists and resource specialists. If you wish to comment or contribute information to this process please contact the MTNHP Botanist. The information we receive from botanists and others throughout the state is essential in this process, and contributes to more accurate assessments of species' status. We continue to ask that all observations for SOC, PSOC and Review Status plants be reported to the Heritage Program. A copy of the field survey form specifying the information that should be submitted is available on our **website** (mtnhp.org).

Information concerning plant species contained on the SOC, PSOC or Review lists may be viewed on the MTNHP's on-line Montana Plant Field Guide. The Field Guide provides information for vascular and non-vascular plants, including species' characteristics, identification, habitat, distribution, state rank reasons and references, as well as technical illustrations and photographs of the plants and their habitats. For each species, a link to the **NatureServe website** (natureserve.org) provides access to information on the status of the species throughout North America, assembled from state and provincial Natural Heritage databases. Information in the Montana Field Guide is continuously updated and expanded, so please check it often for current species' information. If you have questions concerning the field guide or find errors or omissions please contact the MTNHP.

Status lists of SOC plants may be queried on-line by county and/or township; taxonomic group or one of several rank/status criteria. More detailed information or additional assistance can be requested from MTNHP using the Information Request function on our **website**, or by phone, e-mail or mail.

How to Read the Lists

The SOC list is organized alphabetically by scientific name (Genus and specific epithet followed by subspecific epithet if any) within the major groups of Vascular Plants, Bryophytes (Mosses and Liverworts) and Lichens. Vascular plants are further sorted by the subgroups: Ferns and Fern Allies, Gymnosperms (if any), Flowering Plants-Dicots and Flowering Plants-Monocots. The list can also be sorted alphabetically by the common name. Additional scientific names as well as the Family name are included in adjacent columns for each species. The nomenclature and taxonomy for many groups of plants continues to change as new research is conducted and published, and as a result no one nomenclatural reference is followed. Publications and web resources which are most relevant to Montana plants include Vascular Plants of Montana (Dorn 1984), NatureServe Explorer, The USDA PLANTS database, Flora of North America (1993-), Grasses of Montana (Lavin and Seibert 2011) and Flora of the Pacific Northwest (Hitchcock and Cronquist 1973). Additionally, an abundance of scientific literature pertinent to Montana plants is available and indispensable in the process of determining the nomenclature and taxonomic concepts used in this report.

Species that have been added to or deleted from the SOC list due to changes in their global or state rank are reported in separate sections below. These changes are also reflected in the date displayed at the top of the report which shows when an addition or deletion to the list last occurred.

County Distribution

Montana counties of record are listed alphabetically with each species. County records of occurrence are determined directly from mapped species occurrences (SO's) in MTNHP databases. A record of occurrence for a particular county may be based on a historical observation which may no longer be extant. Additionally, some plant observations with vague locality information are not mapped in MTNHP databases and as result would not be included in the county distribution for that particular species.

Montana Species Ranking Codes (GRank, SRank)

Montana employs a standardized ranking system to denote **global** (range-wide) and **state** status (NatureServe 2006). Species are assigned numeric ranks ranging from 1 (highest risk, greatest concern) to 5 (demonstrably secure), reflecting the relative degree of risk to the species' viability, based upon available information.

A number of factors are considered in assigning ranks — the number, size and quality of known occurrences or populations, distribution, trends (if known), intrinsic vulnerability, habitat specificity, and definable threats. The process of assigning state ranks for each taxon relies heavily on the number of occurrences and Species Occurrence (OE) ranks, which is a ranking system of the quality (usually A through D) of each known occurrence based on factors such as size (# of individuals) and habitat quality. The remaining factors noted above are also incorporated into the ranking process when they are known. The "State Rank Reason" field in the **Montana Field Guide** provides additional information on the reasons for a particular species' rank.

Rank Definition

G1 S1	At high risk because of extremely limited and/or rapidly declining population numbers, range and/or habitat, making it highly vulnerable to global extinction or extirpation in the state.
G2 S2	At risk because of very limited and/or potentially declining population numbers, range and/or habitat, making it vulnerable to global extinction or extirpation in the state.
G3 S3	Potentially at risk because of limited and/or declining numbers, range and/or habitat, even though it may be abundant in some areas.
G4 S4	Apparently secure, though it may be quite rare in parts of its range, and/or suspected to be declining.
G5 S5	Common, widespread, and abundant (although it may be rare in parts of its range). Not vulnerable in most of its range.
GX SX	Presumed Extinct or Extirpated - Species is believed to be extinct throughout its range or extirpated in Montana. Not located despite intensive searches of historical sites and other appropriate habitat, and small likelihood that it will ever be rediscovered.
GH SH	Historical, known only from records usually 40 or more years old; may be rediscovered.
GNR SNR	Not Ranked as of yet.
GU SU	Unrankable - Species currently unrankable due to lack of information or due to substantially conflicting information about status or trends.
GNA SNA	A conservation status rank is not applicable because the species or ecosystem is not a suitable target for conservation activities as a result of being: 1) not confidently present in the state; 2) non-native or introduced; 3) a long distance migrant with accidental or irregular stopovers; or 4) a hybrid without conservation value.

Combination or Range Ranks

G#G#

or Indicates a range of uncertainty about the status of the species (*e.g.*, *G1G3 = Global Rank ranges between G1 and G3*).

S#S#

S#, S# Indicates that populations in different geographic portions of the species' range in Montana have a different conservation status (*e.g.*, *S1 west of the Continental Divide and S4 east of the Continental Divide*).

Sub-rank

T# Rank of a subspecies or variety. Appended to the global rank of the full species, *e.g.* *G4T3*

Qualifiers

Q **Questionable** taxonomy that may reduce conservation priority-Distinctiveness of this entity as a taxon at the current level is questionable; resolution of this uncertainty may result in change from a species to a subspecies or hybrid, or inclusion of this taxon in another taxon, with the resulting taxon having a lower-priority (numerically higher) conservation status rank. Appended to the global rank, *e.g.* *G3Q*

? **Inexact Numeric Rank** - Denotes uncertainty; inexactness.

HYB Hybrid - Entity not ranked because it represents an interspecific hybrid and not a species.

C **Captive or Cultivated Only** - Species at present exists only in captivity or cultivation, or as a reintroduced population not yet established.

A **Accidental** - Species is accidental or casual in Montana, in other words, infrequent and outside usual range. Includes species (usually birds or butterflies) recorded once or only a few times at a location. A few of these species may have bred on the few occasions they were recorded.

SYN Synonym - Species reported as occurring in Montana, but the Montana Natural Heritage Program does not recognize the taxon; therefore the species is not assigned a rank.

B **Breeding** - Rank refers to the breeding population of the species in Montana. Appended to the state rank, *e.g.* *S2B, S5N = At risk during breeding season, but common in the winter*

N **Nonbreeding** - Rank refers to the non-breeding population of the species in Montana. Appended to the state rank, *e.g.* *S5B, S2N = Common during breeding season, but at risk in the winter*

M **Migratory** - Species occurs in Montana only during migration.

Federal Status

Designations in this column reflect the status of a species under the U.S. Endangered Species Act (ESA), or as “sensitive” by the U.S. Forest Service (USFS) or Bureau of Land Management (BLM).

U.S. Fish and Wildlife Service (Endangered Species Act)

Status of a taxon under the federal Endangered Species Act of 1973 (16 U.S.C.A. § 1531-1543 (Supp. 1996))

Designation Descriptions	
LE	Listed endangered: Any species in danger of extinction throughout all or a significant portion of its range (16 U.S.C. 1532(6)).
LT	Listed threatened: Any species likely to become an endangered species within the foreseeable future throughout all or a significant portion of its range (16 U.S.C. 1532(20)).
C	Candidate: Those taxa for which sufficient information on biological status and threats exists to propose to list them as threatened or endangered. We encourage their consideration in environmental planning and partnerships; however, none of the substantive or procedural provisions of the Act apply to candidate species.
P	Proposed threatened: Any species that is proposed in the Federal Register to be listed under section 4 of the Act.
DM	Recovered, delisted, and being monitored - Any previously listed species that is now recovered, has been delisted, and is being monitored.
NL	Not listed - No designation.
XE	Experimental - Essential population - An experimental population whose loss would be likely to appreciably reduce the likelihood of the survival of the species in the wild.
XN	Experimental - Nonessential population - An experimental population of a listed species reintroduced into a specific area that receives more flexible management under the Act.
CH	Critical Habitat - The specific areas (i) within the geographic area occupied by a species, at the time it is listed, on which are found those physical or biological features (I) essential to conserve the species and (II) that may require special management considerations or protection; and (ii) specific areas outside the geographic area occupied by the species at the time it is listed upon determination that such areas are essential to conserve the species.
PS	Partial status - status in only a portion of the species' range. Typically indicated in a "full" species record where an infraspecific taxon or population, that has a record in the database has USESA status, but the entire species does not. For example, Yellow-billed Cuckoo (<i>Coccyzus americanus</i>) is ranked PS:LT . Partial Status - Listed Threatened. Designated as Threatened in the Western U.S. Distinct Population Segment (DPS) (subspecies <i>occidentalis</i>)
BGEPA	The Bald and Golden Eagle Protection Act of 1940 (BGEPA) - (16 U.S.C. 668-668c) prohibits anyone, without a permit issued by the Secretary of the Interior, from taking bald or golden eagles, including their parts, nests, or eggs. The BGEPA provides criminal and civil penalties for persons who take, possess, sell, purchase, barter, offer to sell, purchase or barter, transport, export or import, at any time or any manner, any bald eagle ... [or any golden eagle], alive or dead, or any part, nest, or egg thereof. The BGEPA defines take as pursue, shoot, shoot at, poison, wound, kill, capture, trap, collect, molest or disturb. "Disturb" means to agitate or bother a bald or golden eagle to a degree that causes, or is likely to cause, based on the best scientific information available, 1) injury to an eagle, 2) a decrease in its productivity, by substantially interfering with normal breeding, feeding, or sheltering behavior, or 3) nest abandonment, by substantially interfering with normal breeding, feeding, or sheltering behavior. In addition to immediate impacts, this definition also covers impacts that result from human-induced alterations initiated around a previously used nest site during a time when eagles are not present, if, upon the eagles return, such alterations agitate or bother an eagle to a degree that injures an eagle or substantially interferes with normal breeding, feeding, or sheltering habits and causes, or is likely to cause, a loss of productivity or nest abandonment.
MBTA	The Migratory Bird Treaty Act (MBTA) - (16 U.S.C. §§ 703-712, July 3, 1918, as amended 1936, 1960, 1968, 1969, 1974, 1978, 1986 and 1989) implements four treaties that provide for international protection of migratory birds. The statute's language is clear that actions resulting in a "taking" or possession (permanent or temporary) of a protected species, in the absence of a U.S. Fish and Wildlife Service (USFWS) permit or regulatory authorization, are a violation of the MBTA. The MBTA states, "Unless and except as permitted by regulations ... it shall be unlawful at any time, by any means, or in any manner to pursue, hunt, take, capture, kill ... possess, offer for sale, sell ... purchase ... ship, export, import ... transport or cause to be transported ... any migratory bird, any part, nest, or eggs of any such bird [The Act] prohibits the taking, killing, possession, transportation, import and export of migratory birds, their eggs, parts, and nests, except when specifically authorized by the Department of the Interior." The word "take" is defined by regulation as "to pursue, hunt, shoot, wound, kill, trap, capture, or collect, or attempt to pursue, hunt, shoot, wound, kill, trap, capture, or collect." The USFWS maintains a list of species protected by the MBTA at 50 CFR 10.13. This list includes over one thousand species of migratory birds, including eagles and other raptors, waterfowl, shorebirds, seabirds, wading birds, and passerines. The USFWS also maintains a list of species not protected by the MBTA . MBTA does not protect species that are not native to the United States or species groups not explicitly covered under the MBTA; these include species such as the house (English) sparrow, European starling, rock dove (pigeon), Eurasian collared-dove, and non-migratory upland game birds.
BCC	The 1988 amendment to the Fish and Wildlife Conservation Act mandates the U.S. Fish and Wildlife Service to identify species, subspecies, and populations of all migratory nongame birds that, without additional conservation actions, are likely to become candidates for listing under the Endangered Species Act. Birds of Conservation Concern 2008 (BCC 2008) is the most recent effort to carry out this mandate. The overall goal of this report is to accurately identify the migratory and non-migratory bird species (beyond those already designated as federally threatened or endangered) that represent the Service's highest conservation priorities. BCC10, BCC11, and BCC17 designations represent inclusion on the Birds of Conservation Concern list for Bird Conservation Region 10, 11, and 17 in Montana, respectively.

Bureau of Land Management (BLM)

BLM Sensitive Species are defined by the BLM 6840 Manual as native species found on BLM-administered lands for which the BLM has the capability to significantly affect the conservation status of the species through management, and either: (1) there is information that a species has recently undergone, is undergoing, or is predicted to undergo a downward trend such that the viability of the species or a distinct population segment of the species is at risk across all or a significant portion of the species range, or; (2) the species depends on ecological refugia or specialized or unique habitats on BLM-administered lands, and there is evidence that such areas are threatened with alteration such that the continued viability of the species in that area would be at risk.

Designation Descriptions	
Endangered	Denotes species that are listed as Endangered under the Endangered Species Act
Threatened	Denotes species that are listed as Threatened under the Endangered Species Act
Sensitive	Denotes species listed as Sensitive on BLM lands

U.S. Forest Service (USFS)

Designation Descriptions	
Endangered	Listed as Endangered (LE) under the U.S. Endangered Species Act.
Threatened	Listed as Threatened (LT) under the U.S. Endangered Species Act.
Proposed	Any species that is proposed in the Federal Register to be listed under section 4 of the Act.

Candidate	Those taxa for which sufficient information on biological status and threats exists to propose to list them as threatened or endangered. We encourage their consideration in environmental planning and partnerships; however, none of the substantive or procedural provisions of the Act apply to candidate species.
Sensitive	U.S. Forest Service Manual (2670.22) defines Sensitive Species on Forest Service lands as those for which population viability is a concern as evidenced by a significant downward trend in population or a significant downward trend in habitat capacity. These designations were last updated in 2011 and they apply only on USFS-administered lands with land management plans finalized prior to 2017. Sensitive Species designations are being replaced by Species of Conservation Concern designations on individual National Forest as revised land management plans are finalized under the 2012 planning rule.
Species of Conservation Concern	A species, other than federally recognized Threatened, Endangered, Proposed, or Candidate species, that is known to occur in the plan area and for which the regional forester has determined that the best available scientific information indicates substantial concern about the species' capability to persist over the long-term in the plan area (36 CFR 219.9). Species of Conservation Concern replace regional forester Sensitive Species on individual National Forests as revised land management plans are finalized under the 2012 planning rule.

Acknowledgements

We would like to gratefully acknowledge the many people who contributed information on plant species' occurrences and distribution throughout Montana over the years -- those contributions are the building blocks of the MTNHP databases and this publication. We encourage you to continue submitting data for SOC, PSOC and Under Review taxa so that status ranks and this document are as accurate and comprehensive as possible.

Selected References

- Dorn, R.D. 1984. *Vascular Plants of Montana*. Mountain West Publishing, Cheyenne, WY. 276 pp.
- Faber-Langendoen, D., L. Master, J. Nichols, K. Snow, A. Tomaino, R. Bittman, G. Hammerson, B. Heidel, L. Ramsay, and B. Young. 2009. *NatureServe Conservation Status Assessments: Methodology for Assigning Ranks*. NatureServe, Arlington, VA. On-line at natureserve.org/publications/ConsStatusAssess_RankMethodology.pdf
- Flora of North America Editorial Committee, eds. 1993+. *Flora of North America North of Mexico*. 8+ vols. New York and Oxford. On-line at <http://hua.huh.harvard.edu/FNA/> and http://www.efloras.org/flora_page?flora_id=1
- Hitchcock, C.L. and A. Cronquist. 1973. *Flora of the Pacific Northwest*. Univ of Washington Press, Seattle, WA.
- IUCN. 2001. *IUCN Red List Categories and Criteria: Version 3.1*. IUCN Species Survival Commission. IUCN, Gland, Switzerland and Cambridge, UK. 30 pp. On-line at: <http://www.iucn.org>
- Lavin, M. and C. Seibert. 2011. *Grasses of Montana*. MSU Herbarium, Dept of Plant Sciences and Plant Pathology, Montana State University, Bozeman, MT. 100 pp. On-line at: <http://gemini.oscs.montana.edu/~mlavin/herb/mtgrass.pdf>
- Lesica, P., G. Moore, K.M. Peterson, and J.H. Rumely. 1984. *Vascular plants of limited distribution in Montana*. Montana Academy of Science Monograph No. 2.
- Lesica, P. and J.S. Shelly. 1991. *Sensitive, threatened and endangered vascular plants of Montana*. Montana Natural Heritage Program, Montana State Library, Helena, Montana.
- Master, L., D. Faber-Langendoen, R. Bittman, G. Hammerson, B. Heidel, J. Nichols, L. Ramsay, and A. Tomaino. 2009. *NatureServe Conservation Status Assessments: Factors for Assessing Extinction Risk*. NatureServe, Arlington, VA. On-line at http://www.natureserve.org/publications/ConsStatusAssess_StatusFactors.pdf
- NatureServe. *NatureServe Explorer: An on-line encyclopedia of life* [web application]. Version 4.7. Arlington, Virginia. Available: explorer.natureserve.org
- Regan, T.J., L.L. Master and G. A. Hammerson. 2004. *Capturing expert knowledge for threatened species assessments: a case study using NatureServe conservation status ranks*. *Acta Oecologia* 26: 95-107.
- Rollins, R.C. 1993. *The Cruciferae of continental North America: Systematics of the mustard family from the Arctic to Panama*. Stanford University Press, Stanford, California. 976 pp.
- USDA, NRCS. *The PLANTS Database* (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

Contact Information

For questions or comments specific to this publication or for specific plant related questions, please contact:

Andrea Pipp
Program Botanist
apipp@mt.gov
(406) 444-3019

For general questions and botany-related data requests please use the Information Request function on our website (mtnhp.org) or the general MTNHP contact info below.

Montana Natural Heritage Program
P.O. Box 201800
1515 E. 6th Ave.
Helena, MT 59620-1800

Phone: (406) 444-5363
Fax: (406) 444-0581
E-mail: mtnhp@mt.gov

Species of Concern

453 Species

All Records (no filtering)

FERNS AND FERN ALLIES (PTERIDOPHYTA)									
35 SPECIES									
SCIENTIFIC NAME COMMON NAME TAXA SORT	OTHER NAMES	FAMILY (SCIENTIFIC) FAMILY (COMMON)	GLOBAL RANK	STATE RANK	USFWS	USFS	BLM	MNPS THREAT CATEGORY	HABITAT
Asplenium trichomanes- ramosum Limestone Maidenhair Spleenwort	Asplenium viride	Aspleniaceae Spleenwort Family	G5	S3				3	
Species Occurrences verified in these Counties: Carbon, Fergus, Flathead, Glacier, Lake, Lewis and Clark, Pondera, Teton State Rank Reason: S3 SOC: Asplenium trichomanes-ramosum plants are never common, grow in habitat that is limited in Montana, and occur where land management (example: national park, wilderness) provides some protections.									
Botrychium ascendens Upward-lobed Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G3	S3		Sensitive - Known on Forests (KOOT)		4	Various Mesic Sites
Species Occurrences verified in these Counties: State Rank Reason: This moonwort species is documented in Montana primarily from the northwest corner of the state. Almost all observations are on federally-managed lands. Most occurrences are small in size and occupy roadsides or other similarly open or disturbed habitats. As such, it is vulnerable to activities such as weed invasion, weed spraying and road maintenance.									
Botrychium campestre Prairie Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G3G4	S1S2				4	Various Mesic Sites
Species Occurrences verified in these Counties: State Rank Reason: Reported from a very small number of sites in Montana. All occurrences are small with the largest population count at a single site being approximately 2 dozen plants. All known sites are in northwest Montana.									
Botrychium crenulatum Wavy Moonwort	Botrychium dusenii	Ophioglossaceae Adder's-Tongue / Moonworts	G4	S3		Sensitive - Known on Forests (BD, KOOT, LOLO) Species of Conservation Concern on Forests (HLC)		4	Various Mesic Sites
Species Occurrences verified in these Counties: State Rank Reason: This moonwort species is known from numerous observations in western Montana. Most populations are located on either National Forest or State lands. Populations are generally small in size and occupy roadsides or other similarly open or disturbed habitats. As such, it is vulnerable to activities such as weed invasion, weed spraying and road maintenance.									
Botrychium gallicomontanum Frenchman's Bluff Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G2	S1S2				4	Grasslands (Fescue)
Species Occurrences verified in these Counties: State Rank Reason: A globally rare species, recently documented in Montana from Glacier National Park									
Botrychium hesperium Western Moonwort	Botrychium matricariifolium, Botrychium michiganense [in part]	Ophioglossaceae Adder's-Tongue / Moonworts	G4	S3		Sensitive - Known on Forests (BD, KOOT)		4	Various Mesic Sites
Species Occurrences verified in these Counties: State Rank Reason: This moonwort species is known from 25-30 extant sites in western Montana, mostly in Glacier National Park or on National Forest lands. Many sites are poorly documented in terms of population size or are small in size, though several sites have been observed with >100 plants. Many populations occur on roadsides or other similarly open or disturbed habitats. As such, the species is vulnerable to activities such as weed invasion, weed spraying and road maintenance.									
Botrychium lanceolatum Lanceleaf Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G5	S3				3	
Species Occurrences verified in these Counties: State Rank Reason: Reported from approximately two dozen sites. Population levels are poorly documented. As this species was not previously tracked in the state, it may be under-reported.									
Botrychium lineare Linearleaf Moonwort	Slender Moonwort	Ophioglossaceae Adder's-Tongue / Moonworts	G3	S1S2				4	Various Mesic Sites
Species Occurrences verified in these Counties: State Rank Reason: This moonwort species is known to occur in western Montana from 6 locations, 5 of which are on federally-managed lands and the remaining site is located in a tribal wilderness area. However, occurrences are generally small in size and occupy roadsides or other similarly open or disturbed habitats. As such, it is vulnerable to activities such as weed invasion, weed spraying and road maintenance.									
Botrychium michiganense Michigan Moonwort	Botrychium hesperium s.l.	Ophioglossaceae Adder's-Tongue / Moonworts	G3	S2				4	Various Mesic Sites

			Species Occurrences verified in these Counties: State Rank Reason: This species recently has been split from <i>B. hesperium</i>, although it has not yet been formally published (Donald Farrar, Iowa State University). Some of the sites for <i>B. hesperium</i> almost certainly belong here. See <i>B. hesperium</i> for additional information on habitat and characteristics which are very similar. This entity would be included within the concept of <i>B. hesperium</i> as used by the Forest Service on their Sensitive species list.					
Botrychium pallidum Pale Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G3	S1S2			4	Grasslands (Fescue)
			Species Occurrences verified in these Counties: State Rank Reason: Reported from a very small number of sites in Montana. All occurrences are small with the largest population count at a single site being approximately 30 plants. All known sites are in northwest Montana.					
Botrychium paradoxum Peculiar Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G3G4	S3		Sensitive - Known on Forests (BD, KOOT) Sensitive - Suspected on Forests (LOLO) Species of Conservation Concern on Forests (FLAT, HLC)	4	Meadows (Mesic Montane/Subalpine)
			Species Occurrences verified in these Counties: State Rank Reason: This moonwort species is known to occur in western Montana from over two dozen extant occurrences, almost all of which are on federally-managed lands. Many occurrences are small in size and occupy mesic meadows and bunchgrass communities. Potential impacts to these sites include livestock grazing, weed invasion and recreational uses. Though some threats exist to individual occurrences, the species as a whole is not highly threatened by any single or combination of potential impacts in the state.					
Botrychium pedunculatum Stalked Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G3G4	S2		Sensitive - Known on Forests (KOOT) Species of Conservation Concern on Forests (FLAT)	4	Forests (Mesic bottomlands)/Open sites
			Species Occurrences verified in these Counties: State Rank Reason: This moonwort species is known to occur in western Montana from approximately a dozen extant occurrences, almost all of which are on National Forest lands. Many occurrences are small in size and occupy western redcedar forests and roadsides or other similarly open or disturbed habitats. Several site records are based upon specimen collections with no available population data; almost all other sites have population counts with <10 plants observed. One site has been observed with >100 plants. Sites could be negatively impacted by timber harvesting or road-related activities.					
Botrychium pinnatum Northern Moonwort	Botrychium boreale ssp. obtusilobum	Ophioglossaceae Adder's-Tongue / Moonworts	G5	S3			4	
			Species Occurrences verified in these Counties:					
Botrychium simplex Least Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G5	S2			4	
			Species Occurrences verified in these Counties:					
Botrychium sp. (SOC) Moonworts (SOC)		Ophioglossaceae Adder's-Tongue / Moonworts	G1G3	S1S3				
			Species Occurrences verified in these Counties: Deer Lodge, Flathead, Glacier, Granite, Jefferson, Lake, Lewis and Clark, Lincoln, Missoula, Park, Pondera, Powell, Ravalli, Sanders, Sweet Grass, Teton State Rank Reason: This is a general record for <i>Botrychium</i> species tracked by MTNHP. MTNHP tracks and maintains observation data for all <i>Botrychium</i> species in the state excluding <i>B. multifidum</i> and <i>B. virginianum</i> which are fairly common and readily identifiable from all other <i>Botrychium</i>s. Global and State Ranks for this record are placeholders only to allow <i>Botrychium</i> SOC to appear in searches using global and state ranks. For information pertinent to specific <i>Botrychium</i> species, please see the individual species' accounts.					
Botrychium sp. 4 Adnate Moonwort	Botrychium adnatum	Ophioglossaceae Adder's-Tongue / Moonworts	G1?	S1S2			4	Grasslands (Fescue)
			Species Occurrences verified in these Counties: State Rank Reason: A tentatively recognized species that has not been formally published; currently known only from northwest Montana.					
Botrychium spathulatum Spoon-leaf Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G3	S1			4	Forests (Mesic bottomlands)/Open sites
			Species Occurrences verified in these Counties: State Rank Reason: One of the rarest moonwort species in Montana, currently reported from 2 sites in northwest Montana. Population levels at these sites are undocumented.					
Botrychium tunux Moosewort		Ophioglossaceae Adder's-Tongue / Moonworts	G3G4	S1			4	
			Species Occurrences verified in these Counties: State Rank Reason: A globally rare species, recently documented in Montana from Glacier National Park.					

Botrychium yaaxudakeit Yakutat Moonwort		Ophioglossaceae Adder's-Tongue / Moonworts	G3G4	S1			4	Open sites (mesic)
Species Occurrences verified in these Counties: State Rank Reason: A globally rare species, recently documented in Montana from Glacier National Park.								
Cryptogramma cascadenis Cascade Rockbrake		Pteridaceae Maidenhair Fern Family	G5	S3			3	
Species Occurrences verified in these Counties: Flathead, Lincoln, Missoula, Ravalli, Sanders State Rank Reason: <i>Cryptogramma cascadenis</i> is known from 11 locations in western Montana, of which 2 locations are poorly defined and considered historical, 5 locations occur in Wilderness areas, and the remaining 4 locations occur on U.S. Forest Service lands. Although the fern is thought to be undercollected and could be more common, current population and location data is needed to remove this plant from the Species of Concern list.								
Dryopteris cristata Crested Shieldfern		Dryopteridaceae Wood Fern Family	G5	S3		Sensitive - Known on Forests (BRT, KOOT, LOLO) Species of Conservation Concern on Forests (FLAT)	3	Wetland/Riparian
Species Occurrences verified in these Counties: Flathead, Glacier, Lake, Lincoln, Missoula, Ravalli, Sanders State Rank Reason: Rare to uncommon in Montana where it is known from scattered occurrences across the western portion of the state. Most documented occurrences are on National Forest lands, though State Trust Lands and private lands also host significant populations.								
Equisetum palustre Marsh Horsetail		Equisetaceae Horsetails	G5	S3			3	
Species Occurrences verified in these Counties: Beaverhead, Cascade, Chouteau, Fergus, Flathead, Glacier, Granite, Judith Basin, Lake, Lewis and Clark, Lincoln, Madison, Missoula, Phillips, Powell, Ravalli, Sanders State Rank Reason: <i>Equisetum palustre</i> is known from a small number of sites in eight counties of western and central Montana. Plant observations from Beaverhead, Granite, Lewis & Clark, and Chouteau counties need to be confirmed with specimens.								
Equisetum pratense Meadow Horsetail		Equisetaceae Horsetails	G5	S2			3	
Species Occurrences verified in these Counties: Beaverhead, Cascade, Chouteau, Flathead, Granite, Judith Basin, Lake, Lincoln, Meagher, Missoula, Park, Powell, Ravalli, Sweet Grass, Teton State Rank Reason: <i>Equisetum pratense</i> has accurately been identified to occur in a few places within Lake, Powell, and Meagher counties of Montana. Observations in other counties need to be verified because <i>Equisetum pratense</i> can easily be mis-identified. Specimens deposited in herbaria outside of Montana will also need to be examined before it can be demonstrated that this plant is more widely distributed.								
Isoetes echinospora Spiny-spore Quillwort	Isoetes tenella	Isoetaceae Quillworts	G5	S3			3	freshwater lakes
Species Occurrences verified in these Counties: Flathead, Lake, Madison, Missoula, Ravalli, Sanders State Rank Reason: <i>Isoetes echinospora</i> is known from 8 occurrences scattered in western Montana. At one occurrence, the species has been observed in 1940, 1967, and 1998 indicating persistence. However, current survey work is needed to document locations, population sizes, and threats.								
Isoetes howellii Howell's Quillwort		Isoetaceae Quillworts	GNR	S3			3	freshwater lakes
Species Occurrences verified in these Counties: Flathead, Glacier, Lake, Missoula State Rank Reason: <i>Isoetes howellii</i> is known from about 5 locations in Northwestern Montana. Based on limited information threats appear to be minimal, but survey work to document locations, population sizes, and threats is greatly needed.								
Isoetes occidentalis Western Quillwort	Isoetes lacustris var. paupercula	Isoetaceae Quillworts	G4G5	S1			3	freshwater lakes
Species Occurrences verified in these Counties: Flathead, Missoula State Rank Reason: <i>Isoetes occidentalis</i> is known from two locations in northwest Montana. Survey work to identify other locations, document population sizes, and determine threats is greatly needed.								
Lycopodium dendroideum Treelike Clubmoss	Lycopodium obscurum var. dendroideum, Dendrolycopodium dendroideum	Lycopodiaceae Club-moss (Lycopod) Family	G5	S2		Sensitive - Known on Forests (KOOT) Species of Conservation Concern on Forests (HLC)	3	Forests (Mesic valley and montane)
Species Occurrences verified in these Counties: Flathead, Glacier, Lewis and Clark, Lincoln State Rank Reason: Rare in Montana where the species has been documented from only a few sites in the northwest corner of the state. Trend data are unavailable. Known populations do not appear to be immediately threatened by any activities. Populations may be susceptible to negative impacts from fire.								
Lycopodium inundatum Northern Bog Clubmoss	Lycopodiella inundata	Lycopodiaceae Club-moss (Lycopod) Family	G5	S2		Sensitive - Suspected on Forests (KOOT) Species of Conservation Concern on Forests (FLAT)	3	Fens

			Species Occurrences verified in these Counties: Flathead, Missoula State Rank Reason: Rare in Montana where it is known from only a few occurrences in the western portion of the state. Trend data are unavailable. One population may be negatively impacted or extirpated in the future by proposed activities and all populations are susceptible to changes in hydrology.						
Lycopodium lagopus Running-pine	Lycopodium clavatum var. lagopus	Lycopodiaceae Club-moss (Lycopod) Family	G5	S2		Sensitive - Known on Forests (KOOT)		3	Alpine
			Species Occurrences verified in these Counties: Flathead, Glacier, Lincoln State Rank Reason: Rare in Montana. Currently known from two occurrences in the northwest portion of the state. Trend data are unavailable. The known sites do not appear likely to be negatively impacted or threatened from human activity at the current time.						
Marsilea oligospora Pepperwort		Marsileaceae Water-Clover Family	G5	S2				3	
			Species Occurrences verified in these Counties: Lake State Rank Reason: Marsilea oligospora has relatively recently been segregated from Marsilea vestita (FNA 1993). It is quite common around Ninepipes National Wildlife Refuge, but has not been documented elsewhere in Montana. Observation data is greatly needed to further assess its distribution and viability in Montana.						
Ophioglossum pusillum Adder's Tongue	Ophioglossum vulgatum [misapplied]	Ophioglossaceae Adder's-Tongue / Moonworts	G5	S3		Sensitive - Known on Forests (KOOT)		3	Fens, Wet meadows
			Species Occurrences verified in these Counties: Flathead, Lake, Lincoln, Missoula State Rank Reason: Rare in Montana, where it is known from a couple dozen fens and wet meadows in the northwest corner of the state. Its viability in the state generally does not appear to be at risk from any human-caused impacts at this time.						
Phegopteris connectilis Northern Beechfern	Thelypteris phegopteris	Thelypteridaceae Beechfern-Marsh Fern Family	G5	S2S3		Sensitive - Known on Forests (KOOT)		2	Forests (Mesic valley to subalpine)
			Species Occurrences verified in these Counties: Flathead, Glacier, Lincoln, Sanders State Rank Reason: Rare in Montana where it is known from the extreme northwest corner of the state to Glacier National Park. Past timber harvesting likely led to declines in the species' abundance and distribution. Invasive weeds (Orange and Meadow Hawkweeds), proposed mining activity, timber harvesting and fires all have the potential to detrimentally impact the species in the future.						
Polystichum kruckebergii Kruckeberg's Swordfern	Kruckeberg's Hollyfern	Dryopteridaceae Wood Fern Family	G4	S2S3				3	Alpine
			Species Occurrences verified in these Counties: Deer Lodge, Flathead, Gallatin, Lake, Sweet Grass State Rank Reason: Sparsely distributed across western Montana on alpine and subalpine cliffs and talus slopes. Very little data are available for the locations in Montana, though the habitats occupied by the species are not generally impacted by human activities or disturbance. Additional survey and monitoring data are needed.						
Polystichum scopulinum Mountain Swordfern	Mountain Hollyfern	Dryopteridaceae Wood Fern Family	G4	S1S2				3	Rock Crevices
			Species Occurrences verified in these Counties: Ravalli, Sanders State Rank Reason: Only two known locations from western Montana. Very little data are available for the known occurrences. Additional surveys are needed.						
Selaginella selaginoides Northern Spikemoss		Selaginellaceae Spike-mosses	G5	S2S3				3	Wet, mossy soil (montane/subalpine)
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Granite, Madison State Rank Reason: Rare in Montana, where it is known from a few occurrences from the southwest portion of the state. Little survey data are available for known occurrences.						

GYMNOSPERM (CONIFERS)								1 SPECIES	
SCIENTIFIC NAME COMMON NAME TAXA SORT	OTHER NAMES	FAMILY (SCIENTIFIC) FAMILY (COMMON)	GLOBAL RANK	STATE RANK	USFWS	USFS	BLM	MNPS THREAT CATEGORY	HABITAT
Pinus albicaulis Whitebark Pine		Pinaceae Fir / Hemlock / Larch / Pine / Spruce	G3G4	S3	P	Sensitive - Known on Forests (BD, BRT, CG, KOOT, LOLO)	SENSITIVE	2	Subalpine forest, timberline
			Species Occurrences verified in these Counties: Beaverhead, Broadwater, Carbon, Cascade, Deer Lodge, Fergus, Flathead, Gallatin, Glacier, Granite, Jefferson, Judith Basin, Lake, Lewis and Clark, Liberty, Lincoln, Madison, Meagher, Mineral, Missoula, Park, Pondera, Powell, Ravalli, Sanders, Silver Bow, Stillwater, Sweet Grass, Teton, Toole, Wheatland State Rank Reason: Whitebark pine is a common component of subalpine forests and a dominant species of treeline and krummholtz habitats. It occurs in almost all major mountain ranges of western and central Montana. Populations of whitebark pine in Montana and across most of western North America have been severely impacted by past mountain pine beetle outbreaks and by the introduced pathogen, white pine blister rust. The results of which have been major declines in whitebark pine populations across large areas of its range. Additionally, negative impacts associated with encroachment and increased competition from other trees, primarily subalpine fir have occurred as a result of fire suppression in subalpine habitats.						

			Species Occurrences verified in these Counties: Beaverhead, Custer, Prairie State Rank Reason: <i>Astragalus ceramicus</i> variety <i>apus</i> is known only from the upper Snake River Plains of southeast Idaho and adjacent Montana, where it is restricted to the Centennial Valley of Beaverhead County. The disruption of natural disturbance regimes, including fire, ungulate grazing and pocket gopher activity, can lead to dune stabilization, reducing the extent of blowout areas with early successional vegetation, upon which this species depends. Portions of its habitat lie on private or public lands without sensitive species management policies in place.					
Astragalus ceramicus var. filifolius Painted Milkvetch	Pottery Milkvetch	Fabaceae Pea Family	G4T4	S3			2	sandy sites, sand dunes
			Species Occurrences verified in these Counties: Big Horn, Carbon, Carter, Dawson, Powder River, Sheridan State Rank Reason: <i>Astragalus ceramicus</i> variety <i>filifolius</i> is associated with sandy soils of the sandhills and sandstone outcrops in eastern Montana. It is known from about 20 occurrences observed mostly from 1983 to 2000. Some populations occur in State Parks. The Flora of the Great Plains (1986) considered it rare for the region except in the Nebraska sandhill area where it was somewhat common. Based on aging data, limited distribution, and an association to specific habitat types it is considered a Species of Concern. Current data on locations, populations sizes, and threats is greatly needed.					
Astragalus convallarius Lesser Rushy Milkvetch	Astragalus diversifolius [misapplied]	Fabaceae Pea Family	G5	S3		Species of Conservation Concern on Forests (HLC)	2	Grasslands (Intermountain)
			Species Occurrences verified in these Counties: Beaverhead, Broadwater, Jefferson, Lewis and Clark State Rank Reason: The distribution of <i>A. convallarius</i> in Montana is limited to two disjunct localities in the state: the Helena Valley vicinity and an area in extreme southwest Montana in Beaverhead County. The species has been and continues to be negatively impacted by development in the Helena area. Past development in the Helena Valley likely eliminated extensive areas of previously occupied habitat resulting in the more fragmented distribution seen today. The grassland habitats this species occupies are also being invaded by several noxious weeds, particularly in the Helena vicinity. However, the species appears to tolerate some levels of disturbance and degradation of habitat quality. Several large occurrences are presently known and some areas of potentially suitable habitat remain unsurveyed.					
Astragalus geyeri Geyer's Milkvetch		Fabaceae Pea Family	G4	S2			3	Sandy sites
			Species Occurrences verified in these Counties: Carbon, Garfield State Rank Reason: Geyer's milkvetch has a very limited distribution in Montana, primarily limited to Carbon County. Size of the population in Montana is estimated to be in the thousands, but population levels likely fluctuate significantly from year to year. Approximately half the populations occur entirely or partially on federally managed lands.					
Astragalus grayi Gray's Milkvetch		Fabaceae Pea Family	G4?	S2S3			3	Sagebrush-Grassland
			Species Occurrences verified in these Counties: Carbon, Fergus State Rank Reason: Rare in the state. Locally restricted to Carbon and Big Horn counties. Population levels, trends and threats to the species are poorly documented. Additional information is needed for the species within Montana.					
Astragalus lackschewitzii Lackschewitz' Milkvetch	Astragalus molybdenus var. lackschewitzii	Fabaceae Pea Family	G2G3	S2S3		Species of Conservation Concern on Forests (HLC)	3	Alpine
			Species Occurrences verified in these Counties: Pondera, Teton State Rank Reason: Montana endemic restricted to high elevation, gravelly and rocky slopes and ridges. Several of the known occurrences are in designated wilderness and the habitats occupied by the species are not generally subject to human disturbance.					
Astragalus oregonus Wind River Milkvetch		Fabaceae Pea Family	G4?	S2			2	Sandy sites/Sagebrush-Grassland
			Species Occurrences verified in these Counties: Carbon State Rank Reason: Wind River milkvetch is a regional endemic known in Montana only from southern Carbon County. Although populations are relatively large, there are few known occurrences in the state and negative impacts or potential impacts to the species from livestock grazing, ORV use, and extractive industries have been noted.					
Astragalus racemosus Raceme Milkvetch		Fabaceae Pea Family	G5	S2S3			3	Grasslands (Clay soils)
			Species Occurrences verified in these Counties: Carter, Fallon State Rank Reason: Raceme milkvetch occurs near the margin of its range in Montana, where several, mostly small populations have been found in Carter and Fallon counties. Its response to grazing is unknown, however it accumulates selenium and may be toxic to livestock. Accurate population and trend data are lacking.					
Astragalus scaphoides Bitterroot Milkvetch		Fabaceae Pea Family	G3	S3		Sensitive - Known on Forests (BD)	3	Sagebrush-grassland
			Species Occurrences verified in these Counties: Beaverhead, Madison State Rank Reason: Bitterroot milkvetch occurs only in Lemhi County, Idaho and Beaverhead County, Montana. In Montana, the documented occurrences are confined to an area from the Grasshopper Creek drainage south to the Tendoy Mountains. The total number of individual plants has been estimated in the tens of thousands, but occupied habitat is likely less than 700 acres.					
Astragalus terminalis Railhead Milkvetch		Fabaceae Pea Family	G3	S2S3			3	Sagebrush steppe

			Species Occurrences verified in these Counties: Beaverhead, Gallatin, Madison State Rank Reason: <i>Astragalus terminalis</i> is a regional endemic known from southwest Montana, east-central Idaho and northwest Wyoming. In Montana it is documented from Beaverhead County and the Upper Madison River Valley. The species appears to be vulnerable to intensive grazing and competition from noxious weeds, at least in low-elevation areas.						
Athysanus pusillus Sandweed		Brassicaceae Mustards	G5	S1S2		Sensitive - Known on Forests (BRT) Sensitive - Suspected on Forests (LOLO)		2	Rock/talus-Mesic
			Species Occurrences verified in these Counties: Ravalli, Sanders State Rank Reason: Known in Montana from a limited area of the Bitterroot Mountains. Only three occurrences have a large number of individuals and several occurrences have populations of spotted knapweed and/or cheatgrass established. Invasive weeds may threaten the long-term viability of the species in Montana.						
Atriplex truncata Wedge-leaf Saltbush		Amaranthaceae Amaranth (Pigweed) Family	G5	S3				4	Wetland/Riparian
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Jefferson, Lake, Lewis and Clark, Madison, Park, Powell State Rank Reason: Known from two extant occurrences; one in the Centennial Valley and the other near Warm Springs. Also, known historically from four collections in the western half of the state. Additional population and trend data are needed to better evaluate the species' vulnerability.						
Bacopa rotundifolia Roundleaf Water-hyssop		Plantaginaceae Plantain Family	G5	S3?				4	Wetland/Riparian
			Species Occurrences verified in these Counties: Cascade, Fergus, Garfield, Phillips, Powder River, Yellowstone State Rank Reason: A rare species known in Montana from only a few observations in the central and eastern portions of the state. However, the species is widely distributed and appears tolerant of brackish waters as well as some degree of nutrient enrichment. As such, it is unclear to what extent the species' viability is at risk in the state and whether it responds negatively to human-induced impacts to water quality. Additional populations of the species are likely to occur in Montana.						
Balsamorhiza hookeri Hooker's Balsamroot	Balsamorhiza hispida , Balsamorhiza hookeri var. hispida	Asteraceae Aster/Sunflowers	G5	S3				3	Sagebrush-grassland
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge State Rank Reason: Known in Montana only from the vicinity of Monida and within the Mount Haggin WMA.						
Berberis nervosa Longleaf Oregon-grape	Mahonia nervosa	Berberidaceae Barberries	G5	S1				4	
			Species Occurrences verified in these Counties: Sanders State Rank Reason: <i>Berberis nervosa</i> is disjunct in northern Idaho. In Montana it is known from 2-3 locations in Sanders County, of which one population in 2001 is reported to have over 1,000 plants. Additional data on locations and population sizes are greatly needed.						
Bidens beckii Beck Water-marigold	Megalodonta beckii	Asteraceae Aster/Sunflowers	G5	S2		Sensitive - Known on Forests (KOOT, LOLO)		3	Aquatic
			Species Occurrences verified in these Counties: Broadwater, Flathead, Lake, Lincoln, Missoula, Powell, Ravalli State Rank Reason: Known from ten occurrences in the western valleys of the state, including 6 moderate to large populations and one historical occurrence from Salmon Lake dating to 1937. However, the species may be more abundant in the state than what current data suggests. Threats and impacts to populations in Montana include boating activity, lake shore development, aquatic weeds and use of aquatic herbicides.						
Boechera demissa Daggett Rockcress	Arabis demissa	Brassicaceae Mustards	G5	S1S3				3	Open woodland and sagebrush steppe
			Species Occurrences verified in these Counties: Carbon State Rank Reason: Daggett rockcress is at the northern edge of its range in Montana, where it is known only from the vicinity of the Pryor Mountains and adjacent Bighorn Canyon. Detailed survey information for most occurrences is lacking.						
Boechera fecunda Sapphire Rockcress	Arabis fecunda	Brassicaceae Mustards	G2	S2		Sensitive - Known on Forests (BD) Sensitive - Suspected on Forests (BRT, LOLO)	SENSITIVE	1	Rocky, calcareous, montane slopes
			Species Occurrences verified in these Counties: Beaverhead, Ravalli, Silver Bow State Rank Reason: Sapphire rockcress is a state endemic known from several locations in southwest Montana where it is restricted to specific and localized habitats. Encroachment of spotted knapweed threatens several populations, particularly in Ravalli County. It is unclear whether grazing has significant negative impacts						
Brasenia schreberi Watershield		Cabombaceae Watershields	G5	S1S2		Sensitive - Known on Forests (KOOT, LOLO)		3	Aquatic
			Species Occurrences verified in these Counties: Flathead, Lake, Lincoln, Missoula, Powell State Rank Reason: Restricted in Montana to shallow waters in the valleys of the northwest corner of the state where it is known from eight occurrences, including six relatively high quality populations. Potential threats to the species include boating activity, aquatic weeds, and several populations are subject to runoff from adjacent agricultural fields, though it is uncertain if this has negatively impacted any populations.						

Braya humilis Low Braya	Neotorularia humilis	Brassicaceae Mustards	G5	S2		Species of Conservation Concern on Forests (HLC)	3	Alpine
Species Occurrences verified in these Counties: Beaverhead, Fergus, Teton State Rank Reason: Known from four locations in the state, including one site in which only one plant was observed. One population occurs in an area with historical mining activity and may have been detrimentally impacted. Another populations occurs along the Rocky Mtn Front and is actively monitored; population levels may be declining at this site based upon preliminary data.								
Brickellia oblongifolia Mojave Brickellbush		Asteraceae Aster/Sunflowers	G5	S1S2			3	Rock/Talus
Species Occurrences verified in these Counties: Park, Silver Bow State Rank Reason: Few collections known for Montana. Only known extant occurrences are all near Melrose. The current status of one historical occurrence near Wilsall is unknown. Invasive weeds do not appear to be a threat at this time and the rocky, sparsely-vegetated slopes that the species occupies are not generally subject to human impacts. Livestock grazing may be negatively impacting the species at one site. Updated population and site data are needed for the known occurrences. Other occurrences of the species are likely to be found in Montana.								
Camissonia andina Obscure Evening-primrose	Oenothera andina, Holmgrenia andina	Onagraceae Evening-primrose Family	G4	S2			3	Sandy sites
Species Occurrences verified in these Counties: Carbon, Missoula State Rank Reason: This species is at the edge of its range in Montana, where it has been documented from just a few locations. All known extant locations are from Carbon County. These populations collectively cover less than 20 acres, but they can vary greatly in size from year to year. It tolerates grazing well, and moderate grazing may be important in maintaining a suitable seedbed of exposed soil. Invasive weeds may pose the greatest risk.								
Camissonia parvula Small Camissonia	Oenothera parvula	Onagraceae Evening-primrose Family	G5	S1S2			4	Sandy sites
Species Occurrences verified in these Counties: Carbon State Rank Reason: <i>Camissonia parvula</i> is currently known from one extant location in Montana on the southern edge of the Pryor Mountains in Carbon County. Populations are thought to be small, but may vary widely from year to year. As an annual plant, it may tolerate - or even respond positively to - moderate levels of disturbance. Additional population and site data are needed for this species in Montana.								
Cardamine oligosperma var. kamtschatica Few-seeded Bittercress	Cardamine umbellata	Brassicaceae Mustards	G5T5	S2?			3	Alpine
Species Occurrences verified in these Counties: Flathead State Rank Reason: Only known from 1 collection in Montana. Additional data are needed to reliably determine the species' conservation status and needs in Montana.								
Cardamine rupicola Cliff Toothwort		Brassicaceae Mustards	G3	S3			3	Alpine
Species Occurrences verified in these Counties: Flathead, Lake, Lewis and Clark, Missoula, Powell State Rank Reason: State endemic known from 3 population clusters. These are in the Mission Mtns, Swan Range and the Rocky Mtn Front Range. Many occurrences have not been surveyed for 30 or more years and many are based on a single herbarium specimen. However, the species grows at high elevations in rock and scree fields that generally are not subject to disturbance or other threats. Many populations also occur in designated wilderness areas which offer further protection. Additional occurrences likely exist across the known range of the species.								
Castilleja covilleana Coville Indian Paintbrush		Orobanchaceae Broomrape Family	G3G4	S3		Sensitive - Known on Forests (BRT) Sensitive - Suspected on Forests (BD)	2	Subalpine slopes
Species Occurrences verified in these Counties: Beaverhead, Flathead, Lake, Missoula, Ravalli State Rank Reason: This species is known in Montana, primarily from the West Fork of the Bitterroot River on the Bitterroot National Forest. 5 occurrences are known from historical collections or have unknown status. A few occurrences contain minor amounts of spotted knapweed and others occur in habitats that are susceptible to invasion by knapweed and other invasive species. Timber harvest activities may also pose a threat to some populations.								
Castilleja exilis Annual Indian Paintbrush	Castilleja minor ssp. minor	Orobanchaceae Broomrape Family	G5T5	S2			3	Wetland/Riparian
Species Occurrences verified in these Counties: Broadwater, Deer Lodge, Fergus, Gallatin, Jefferson, Madison, Park State Rank Reason: Annual Indian Paintbrush is known from a half dozen counties in southwest Montana with the majority of documented locations on private lands. Many areas of suitable habitat have been converted to agricultural uses and/or are used for livestock grazing. Additionally, populations are susceptible to hydrologic changes and may be negatively impacted by invasive weeds.								
Castilleja gracillima Slender Indian Paintbrush	Castilleja miniata ssp. miniata	Orobanchaceae Broomrape Family	G3G4	S2			3	Wetland/Riparian
Species Occurrences verified in these Counties: Beaverhead, Gallatin, Madison, Park State Rank Reason: This plant is a regional endemic, known in Montana from a limited number of populations, with most being relatively small. No threats have been observed, though it could be vulnerable to hydrologic alterations or noxious weeds.								
Castilleja kerryana Kerry's Paintbrush		Orobanchaceae Broomrape Family	G3	S3		Species of Conservation Concern on Forests (HLC)	3	

			Species Occurrences verified in these Counties: Lewis and Clark State Rank Reason: Castilleja kerryana is a recently recognized species that grows in alpine habitat in a portion of the Scapegoat Wilderness in Montana. Populations tend to be small and scattered on slopes and ridges, and apparently absent on broad, fairly flat alpine terrain. Although Castilleja species in general have brittle stems that are easily damaged by livestock, grazing is not known to occur where Kerry's Paintbrush grows. The plant appears to be limited geographically in Montana, and additional surveys are needed to accurately determine its range.					
Castilleja nivea Snow Indian Paintbrush		Orobanchaceae Broomrape Family	G3	S3			3	Alpine
			Species Occurrences verified in these Counties: Carbon, Fergus, Golden Valley, Madison, Park, Sweet Grass State Rank Reason: Currently known from a few collections from the Beartooths, Crazy Mtns, Tobacco Root Mtns and the Centennial Range. It is very likely that additional occurrences exist in the known mountain ranges as well as additional mountain ranges. Additionally, the high elevation habitat generally limits the potential for impacts to the species.					
Celastrus scandens Bittersweet		Celastraceae Bittersweet Family	G5	S1			2	Wetland/Riparian
			Species Occurrences verified in these Counties: Carter, Dawson, Richland State Rank Reason: Celastrus scandens occurs frequently in woodlands, rocky hillsides, thickets, fence rows, and roadsides in the Great Plains (McGregor et al. 1986). The previous Montana rank of SH was based on a vague location provided on a 1975 herbarium specimen. In recent years it has been collected at four locations in woody draws. It appears that the Montana sites represent the western edge of its range, and currently it ranks as an S1. Additional surveys of woody draws are needed to accurately document its distribution and population size in Montana.					
Centunculus minimus Chaffweed	Anagallis minima, Lysimachia minima	Myrsinaceae Myrsine Family	G5	S2			3	Wetland/Riparian
			Species Occurrences verified in these Counties: Cascade, Lake, Missoula, Phillips, Powell, Ravalli, Sheridan, Valley State Rank Reason: Known from scattered locations across the state, though it is rare to uncommon in Montana. May be susceptible to some adverse impacts from human-caused disturbance due to its preference for vernal moist habitats in valley loctions.					
Cercocarpus montanus Alderleaf mountain-mahogany		Rosaceae Rose Family	G5	S2S3			3	Open, stony slopes
			Species Occurrences verified in these Counties: Dawson, Treasure State Rank Reason: This widespread western species is only known in the state from one area of Treasure County where it is reported to be fairly extensive. Additional data on population size and extent are needed to more precisely rank the species.					
Chenopodium subglabrum Smooth Goosefoot	Chenopodium leptophyllum var. subglabrum	Amaranthaceae Amaranth (Pigweed) Family	G3G4	S2			2	Sandy sites
			Species Occurrences verified in these Counties: Carter, Cascade, Custer, Fergus, Garfield, Mccone, Phillips, Powder River, Sheridan State Rank Reason: Smooth goosefoot is known from just a few locations in Montana, one of which may be extirpated. It occupies an early-succession habitat that is vulnerable to loss of natural disturbance regimes such as fire and flooding. Invasion of exotic plants may also pose a threat. Population data and trend monitoring data are lacking though the populations likely fluctuate widely from year to year.					
Cirsium longistylum Long-styled Thistle		Asteraceae Aster/Sunflowers	G2G3	S2S3			2	Meadows (Montane-subalpine)
			Species Occurrences verified in these Counties: Broadwater, Cascade, Fergus, Judith Basin, Lewis and Clark, Meagher, Wheatland State Rank Reason: Population estimates of approximately 30,000 plants, including seven high quality populations, scattered over four mountain ranges are promising for the long-term viability of the species. Habitat in the largest populations is generally of high quality with few if any problem weeds posing significant and immediate threats. In the near future, little change in habitat quality is expected in these populations. Sites are mostly on National Forest lands that provide a degree of protection and two large populations on private lands that have a history of light to moderate grazing appear stable. Also of benefit at this time is the active weed control program employed by the private landowners on their lands. Long- and short-term population trends are difficult to gauge due to the lack of good survey data over many years. However, available data and observations provide some evidence that population levels have at least remained fairly stable over the past decade, with significant yearly fluctuations possible. Threats posed by invasive weeds and the introduced bio-control agent do provide reason for concern.					
Cirsium pulcherrimum Wyoming Thistle		Asteraceae Aster/Sunflowers	G5	S3			3	Sparsely-vegetated soils
			Species Occurrences verified in these Counties: Big Horn, Carbon, Powder River, Prairie State Rank Reason: Known in Montana from one badlands area of Powder River County with a small number of scattered individuals observed in 2006. Also, reported for Dawson and Garfield Counties by Flora of the Great Plains and 1 collection from each of Carbon and Custer Counties.					
Clarkia rhomboidea Diamond Clarkia		Onagraceae Evening-primrose Family	G5	S3		Sensitive - Known on Forests (BRT, KOOT, LOLO)	2	Forests (Open, montane)
			Species Occurrences verified in these Counties: Lake, Lincoln, Ravalli, Sanders State Rank Reason: Rare in Montana, where it is known from only a small portion of the northwest corner of the state, primarily along the lower Clark Fork River drainage. Some detrimental impacts from invasive weeds and subsequent herbicide treatments are possible as are loss of habitat due to fire suppression.					
Claytonia arenicola Sand Springbeauty	Montia arenicola	Portulacaceae Purslane Family	G4	S2S3		Sensitive - Known on Forests (LOLO)	3	Mesic, rocky slopes
			Species Occurrences verified in these Counties: Sanders State Rank Reason: Rare in Montana, where it is currently known from only one localized area in the western portion of the state. As an annual, populations likely fluctuate widely from year to year. No specific threats have been identified.					
Cleome lutea Yellow Beeplant	Peritoma lutea	Cleomaceae Cleome Family	G5	S1S2			3	Sagebrush-grassland (Low-elevation)

			Species Occurrences verified in these Counties: Big Horn, Carbon, Deer Lodge State Rank Reason: Rare in Montana, where it is currently known from only a small area in the south-central portion of the state. Current population levels and trends are undocumented, though populations likely fluctuate widely from year to year. Additional monitoring is needed.						
Collomia debilis var. camporum Alpine Collomia		Polemoniaceae Phlox Family	G5T2	S1S2				3	Rock/Talus (Valleys to Montane)
Species Occurrences verified in these Counties: Granite, Missoula, Ravalli State Rank Reason: Only known from a few sites in western Montana and Lemhi County, Idaho, from low elevation scree, talus or rocky slopes. Negative impacts from human disturbance and weed invasion are possible. Current status of most of the documented locations is not known. Survey and monitoring data are needed.									
Corydalis sempervirens Pale Corydalis	Capnoides sempervirens	Fumariaceae Fumary family	G5	S2		Sensitive - Known on Forests (KOOT) Species of Conservation Concern on Forests (FLAT)		3	Forests/Meadows (Recently-burned)
Species Occurrences verified in these Counties: Flathead, Glacier, Lincoln, Powell State Rank Reason: Known to occur in northwest Montana from approximately a dozen recently documented (past 25 years) occurrences. Another five historical occurrences are also known. This species occurs in disturbed habitats, predominantly burned forests and it depends heavily on historical fire regimes to maintain populations. Thus, the main threat to this species' viability appears to be from fire suppression activities. Invasive weeds also threaten habitat occupied by the species.									
Cryptantha fendleri Fendler Cat's-eye		Boraginaceae Borage Family	G5	S2			SENSITIVE	2	Sandy sites
Species Occurrences verified in these Counties: Beaverhead, Gallatin, Sheridan State Rank Reason: Fendler cat's-eye is restricted to very localized sandhills habitat in the far southwestern and northeastern corners of Montana where it is known from a total of three moderate to large-sized populations. It responds positively to disturbance that maintains its sparsely vegetated habitat. Fire suppression and dune stabilization efforts have likely had an adverse effect on populations of this species.									
Cryptantha humilis Round-headed Cryptantha		Boraginaceae Borage Family	G4?	SH				4	Sagebrush Steppe (low-elevation)
Species Occurrences verified in these Counties: Beaverhead, Jefferson State Rank Reason: Known from 3 historical collections in the state, including a 1955 collection west of Dillon in the Grasshopper Valley, a 1952 collection 3 miles south of Lima and an undated collection from the Yellowstone Valley in Park County.									
Cryptantha scoparia Miner's Candle		Boraginaceae Borage Family	G4?	S2				4	Sagebrush Steppe (low-elevation)
Species Occurrences verified in these Counties: Carbon State Rank Reason: This species is documented from a single area in Carbon County, where it is widely disjunct from the nearest known occurrences in southwest Wyoming and central Idaho. In 1991 about 1,000 plants were reported occupying less than one acre. The habitat is subject to grazing, and may be affected by exotic weed encroachment. Additional surveys and monitoring data are needed.									
Dalea enneandra Nine-anther prairie clover		Fabaceae Pea Family	G5	S2S3				3	Grasslands (Plains)
Species Occurrences verified in these Counties: Big Horn, Custer, Fallon, Richland State Rank Reason: In Montana, known from a few poorly documented occurrences in the eastern half of the state. Additional surveys and updated population data are needed.									
Dalea villosa Silky prairie clover	Petalostemon villosus	Fabaceae Pea Family	G5	S2				4	Sandy sites
Species Occurrences verified in these Counties: Carter, Fallon, Richland, Sheridan State Rank Reason: In Montana, known from a few, small occurrences in the extreme eastern portion of the state. Current population levels and trends are unknown.									
Delphinium burkei Meadow Larkspur	[including] Delphinium distichum	Ranunculaceae Buttercup Family	G4	S1S2				4	Meadows (Moist, low-elevation)
Species Occurrences verified in these Counties: Beaverhead, Flathead, Silver Bow State Rank Reason: Only known from a few collections from the western half of the state.									
Delphinium depauperatum Slim Larkspur		Ranunculaceae Buttercup Family	G5	S2				3	
Species Occurrences verified in these Counties: Beaverhead, Flathead, Madison, Pondera State Rank Reason: Delphinium depauperatum has been identified in Beaverhead, Flathead, and possibly Jefferson Counties in western Montana. It is found in common habitats, yet relatively few occurrences have been documented.									
Delphinium glaucum Pale Larkspur		Ranunculaceae Buttercup Family	G5	S1?				3	
Species Occurrences verified in these Counties: Beaverhead, Gallatin, Madison, Mineral, Park, Stillwater State Rank Reason: Based on the discrepancy in the number of herbarium specimens identified as Delphinium glaucum (CPNWH 2015) and in its Montana County distribution (Lesica 2012), there seems to be an issue in how to accurately identify this species. Specimens deposited in herbaria outside of Montana will need to be examined before it can be demonstrated that this plant is more widely distributed.									

Drosera anglica English Sundew		Droseraceae Sundew Family	G5	S3		Sensitive - Known on Forests (BD, BRT, CG, KOOT, LOLO) Species of Conservation Concern on Forests (HLC)		3	Fens
			Species Occurrences verified in these Counties: Beaverhead, Flathead, Granite, Lake, Lewis and Clark, Lincoln, Madison, Missoula, Park, Powell, Ravalli, Sanders State Rank Reason: Known from over two dozen populations in the state, most of these are moderate to large-sized, healthy populations. Most occurrences are on federally managed lands with several of these in designated wilderness areas, research natural areas or Glacier National Park which help to protect the occurrences from many potential threats. However, one population is vulnerable to ski area expansion and activity, and the species may be negatively impacted by fire as observations at one location appear to indicate. Plants are also sensitive to and negatively impacted by trampling of peat mats on which the species grow.						
Drosera linearis Slenderleaf Sundew		Droseraceae Sundew Family	G4G5	S2		Sensitive - Suspected on Forests (KOOT) Species of Conservation Concern on Forests (FLAT, HLC)		3	Fens
			Species Occurrences verified in these Counties: Flathead, Lake, Lewis and Clark, Powell State Rank Reason: Only known from four populations in Montana though all are moderate to large-sized occurrences that are located in either the Bob Marshall Wilderness or Indian Meadows Research Natural Area which afford all known populations some protection from disturbance.						
Dryas integrifolia Entire-leaved Avens		Rosaceae Rose Family	G5	S2S3				4	Alpine
			Species Occurrences verified in these Counties: Fergus, Golden Valley State Rank Reason: Known in Montana from the Big Snowy Mountains and possibly from the Tobacco Root Mountains, though location of this latter specimen collection is unknown and cannot be confirmed. Current population levels and trends are unknown. However, its high-elevation habitat is relatively inaccessible, and there does not appear to be any significant threats.						
Ericameria discoidea var. discoidea Whitestem Goldenbush	Haplopappus macronema var. macronema	Asteraceae Aster/Sunflowers	G4	S2		Sensitive - Known on Forests (BD, CG) Sensitive - Suspected on Forests (BRT)		3	Rock/Talus
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: Rare in Montana where it is only known from a couple of sites in the southwest corner of the state. Population levels are poorly documented. One site is relatively inaccessible and not likely to be threatened by human impacts.						
Ericameria parryi var. montana Parry's Mountain Rabbitbrush	Chrysothamnus parryi ssp. montanus	Asteraceae Aster/Sunflowers	G5T2	S2				3	Grasslands (subalpine)
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: A globally rare endemic, restricted to a small area of southwest Montana and adjacent Idaho. Though only known from one population in Montana with an estimated couple hundred plants, its habitat is remote and there are no apparent threats to its viability in the near future. Additional data on population levels and trend should be collected.						
Erigeron allocotus Big Horn Fleabane		Asteraceae Aster/Sunflowers	G3	S3			SENSITIVE	3	Rock outcrops/Ridges (low-elevation)
			Species Occurrences verified in these Counties: Big Horn, Carbon State Rank Reason: A regional endemic of Montana and Wyoming. In Montana, it is known only from the Pryor Mountain Desert - Bighorn Basin area of Carbon and Big Horn Counties. The species can be common in areas where it is found.						
Erigeron asperugineus Idaho Fleabane		Asteraceae Aster/Sunflowers	G4	S2		Sensitive - Known on Forests (BD, BRT)		3	Alpine
			Species Occurrences verified in these Counties: Beaverhead, Ravalli State Rank Reason: Idaho fleabane is a regional endemic that has been documented from a few locations in Montana. It grows in alpine habitats, which tend to be relatively isolated from anthropogenic disturbance. Updated population data are needed for most occurrences and it is likely that a few additional occurrences will be documented.						
Erigeron evermannii Evermann Fleabane		Asteraceae Aster/Sunflowers	G4	S2?		Sensitive - Known on Forests (BRT)		3	Alpine
			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Rare in Montana, where it is currently known from two alpine peaks in the Bitterroot Mountains. Available data are based on specimen collections from the 1960's and 1970's, though there is no reason to believe that these populations no longer exist or that they have been negatively impacted. More current data are needed.						

Grindelia howellii Howell's Gumweed	Grindelia paysonorum	Asteraceae Aster/Sunflowers	G3	S2S3		Sensitive - Known on Forests (LOLO) Sensitive - Suspected on Forests (KOOT) Species of Conservation Concern on Forests (FLAT, HLC)	SENSITIVE	2	Vernally moist sites (Open, Low-elevation)
Species Occurrences verified in these Counties: Granite, Missoula, Powell State Rank Reason: In Montana, <i>Grindelia howellii</i> is known from over 100 mapped occurrences. However, most populations are small and many occur on roadsides or other similarly disturbed habitat. This habitat preference in conjunction with the short-lived nature of the species means occurrences may drift from place to place or from year to year and as a result many occurrences may be ephemeral. These attributes make determination of population numbers as well as the number of extant populations at any given time difficult to assess. Invasive weeds are a threat to many occurrences, as the habitat occupied by <i>G. howellii</i> is also favorable for many weedy species. Application of herbicides to control these weeds, especially along roadsides may also have a direct, negative impact.									
Gymnosteris parvula Small-flower Gymnosteris		Polemoniaceae Phlox Family	G4	S2				3	Grasslands/Sagebrush steppe
Species Occurrences verified in these Counties: Beaverhead, Gallatin State Rank Reason: Known in Montana from one 1932 collection near West Yellowstone and one recent collection from Beaverhead County.									
Helianthus pumilus Little Sunflower		Asteraceae Aster/Sunflowers	G4?	S1				4	
Species Occurrences verified in these Counties: Big Horn State Rank Reason: <i>Helianthus pumilus</i> is documented from a single 2007 collection from Big Horn County, Montana (Rocky Mountain Herbarium #910246). It was previously reported for Montana by the <i>Flora of the Great Plains</i> (McGregor et al. 1986). Globally this plant occurs in Colorado and Wyoming where it occupies dry, rocky places (Schilling <i>in</i> Flora of North America 2007). This location, though imprecisely mapped, extends the range northward. Surveys that bring forth precise mapping and data on population sizes, habitat, and threats are greatly needed.									
Heterocodon rariflorum Western Pearl-flower		Campanulaceae Bellflower Family	G5	S2		Sensitive - Known on Forests (BRT, KOOT, LOLO)		3	Vernally moist habitats
Species Occurrences verified in these Counties: Beaverhead, Lake, Lincoln, Mineral, Missoula, Powell, Ravalli, Sanders State Rank Reason: Over a dozen known occurrences, including a half-dozen moderate to large-sized populations, a few small populations and several occurrences that need further survey work to document population sizes. Most populations are on National Forest lands. Invasive weeds infest several populations and are likely infest others. Hiking and ORV trails occur though or adjacent to a few populations and associated use may impact <i>H. rariflorum</i> plants.									
Hornungia procumbens Hutchinsia	Hutchinsia procumbens	Brassicaceae Mustards	G5	S2				3	Sagebrush Steppe
Species Occurrences verified in these Counties: Beaverhead, Carbon, Flathead, Powell State Rank Reason: Rare in Montana. Currently known from approximately a half-dozen occurrences scattered across the mountainous portion of the state. Trend and population data are generally lacking, though it is an annual and populations probably fluctuate widely from year to year. Threats to the species' viability in Montana appear to minimal.									
Howellia aquatilis Water Howellia		Campanulaceae Bellflower Family	G3	S3	DM			2	Aquatic
Species Occurrences verified in these Counties: Lake, Missoula State Rank Reason: Water howellia is restricted in Montana to depressional wetlands in the Swan Valley, typically occupying small basins where the water level recedes partially or completely by the Fall. Montana contains the largest number of occupied ponds and wetlands though the total occupied area is small and it is clustered in a small portion of the state, making it vulnerable to localized events and management actions. Reed canary grass (<i>Phalaris arundinacea</i>) has invaded into some wetlands in the Swan Valley and it has the potential to form dense monocultures, thereby decreasing the amount of available habitat, though it has only been found in a small percentage of occupied water howellia sites so far. Additionally, water howellia is an annual species, which is solely dependent on recruitment from seed; and it has very narrow habitat and moisture requirements which leaves it vulnerable to extirpation as a result of consecutive years of unfavorable growing conditions.									
Idahoia scapigera Scalegod		Brassicaceae Mustards	G5	S1S2		Sensitive - Known on Forests (BRT) Sensitive - Suspected on Forests (LOLO) Species of Conservation Concern on Forests (FLAT)		2	Vernally moist, rock ledges

			Species Occurrences verified in these Counties: Flathead, Ravalli, Sanders State Rank Reason: Rare and peripheral in Montana. Currently known from approximately a half-dozen sites in western Montana, mostly along the lower slopes of the Bitterroot Mountains. Populations are highly susceptible to negative impacts from invasive weeds, primarily spotted knapweed and cheatgrass. Data on population trends are lacking, though levels likely fluctuate widely from year to year.					
Impatiens aurella Pale-yellow Jewel-weed		Balsaminaceae Impatiens	G4	S3			3	riparian
			Species Occurrences verified in these Counties: Cascade, Flathead, Gallatin, Jefferson, Lake, Lewis and Clark, Mineral, Missoula, Sanders State Rank Reason: <i>Impatiens aurella</i> is known from about 20 locations documented from 1886 to 2016. It is considered uncommon in Lake and Flathead Counties, where the majority of observations have been found, and rare in other counties of western Montana. It grows in wet, often organic soil in both disturbed and undisturbed wetlands, and rarely appears abundant. However, it may require or persist better with some hydrological disturbance. Revisits to known locations and more surveys are needed to better document locations, population sizes, and threats.					
Ipomoea leptophylla Bush morning-glory		Convolvulaceae Morning-glory Family	G3G5	S1S2			4	Prairie
			Species Occurrences verified in these Counties: Big Horn, Rosebud, Treasure, Yellowstone State Rank Reason: Known in Montana from only a few collections in the southeastern part of the state, only 1 of these collections was in the last 2 decades. This is a very conspicuous, attractive species, so it is probably not undercollected.					
Ipomopsis congesta ssp. crebrifolia Ballhead Ipomopsis	Gilia congesta var. crebrifolia Ballhead Gilia, Ball-head Standing-cypress, Compact Gilia	Polemoniaceae Phlox Family	G5T3T4	S2S3			3	Sagebrush Steppe
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: Rare and peripheral in Montana. Currently known from only a small geographic area encompassing parts of the Centennial Mountains to the Monida Pass area in southwest Montana. Additional data on population levels are needed, though it is expected that populations are stable. Potential threats to the known occurrences appear to be minimal or non-existent at the current time.					
Ipomopsis minutiflora Small-flower Ipomopsis	Gilia minutiflora, Microgilia minutiflora Small-flower Standing-cypress	Polemoniaceae Phlox Family	G4	S1S2			4	Sagebrush (Open)
			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Rare and peripheral in Montana. Currently documented in the state from one collection from the Bitterroot Valley. Very little is known about this species in the state. Additional surveys are needed. Species may be overlooked/undercollected or perhaps the Montana occurrence could be the result of a more recent and isolated establishment event.					
Kelloggia galioides Kelloggia		Rubiaceae Bedstraws / Madder Family	G5	SH			4	Forest (Open/low-elevation)
			Species Occurrences verified in these Counties: Mineral State Rank Reason: Known in Montana from one 1971 collection in the South Fork Fish Creek valley approximately 12 miles west-northwest of Alberton and a 0.5 mile north of the junction with Deer Creek.					
Kochia americana Red Sage	Bassia americana Green Molly	Amaranthaceae Amaranth (Pigweed) Family	G5	S2			2	Saline/Alkaline Sites
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: The species is at the periphery of its range in Beaverhead County where it is known from one large extant population on BLM and private lands, two historical locations and two other locations that need additional survey work. Agricultural conversion has significantly reduced available habitat. Additional impacts to <i>K. americana</i> from agriculture, grazing and/or invasive weeds are possible.					
Koenigia islandica Island Koenigia		Polygonaceae Buckwheat Family	G4	S2			3	Alpine
			Species Occurrences verified in these Counties: Carbon, Flathead, Sanders State Rank Reason: Rare in Montana, where it is only known from several, high elevation sites on the Beartooth Plateau. Data are insufficient for accurately determining population levels and trend, though populations probably fluctuate widely from year to year. The known occurrences and their habitat do not appear to be at any significant risk of adverse impacts from human activities.					
Lagophylla ramosissima Slender Hareleaf		Asteraceae Aster/Sunflowers	G5	S1			2	Grasslands (Dry/Valley)
			Species Occurrences verified in these Counties: Lake, Sanders State Rank Reason: Species is poorly documented in Montana where it is known from three occurrences in close proximity to each other. More survey work for the species is needed to determine sizes of existing populations at a minimum. Invasive weeds occur at or near existing sites, though impacts of invasive weeds on <i>L. ramosissima</i> are unknown.					
Lathyrus bijugatus Latah Tule Pea		Fabaceae Pea Family	G4	S2S3		Sensitive - Known on Forests (KOOT)	3	Forest (Open/Valley)
			Species Occurrences verified in these Counties: Flathead, Lincoln State Rank Reason: Rare and peripheral in Montana. Currently documented from three, widely scattered sites in the valleys-lower mountains of northwest Montana.					
Leptodactylon caespitosum Matted Prickly-phlox	Linanthus caespitosus, Linanthus cespitosus	Polemoniaceae Phlox Family	G4	S2S3			2	Sandy Breaks/Outcrops
			Species Occurrences verified in these Counties: Carbon State Rank Reason: This plant occurs in Montana at the edge of a broad but patchy range. It is known from only a dozen or so mostly small populations, all in the Pryor Mountains - Bighorn Canyon area, and is confined to a highly specific substrate. The habitat of this plant receives little human disturbance and there are no evident threats.					
Lewisia columbiana Columbia Lewisia		Portulacaceae Purslane Family	G4G5	S1S2			1	Rock Crevices

			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Rare and peripheral in Montana, where it is known from only one location in the Bitterroot Mountains. Its relatively inaccessible habitat reduces the potential for negative impacts.					
Ligusticum verticillatum Idaho Lovage		Apiaceae Parsley/Carrot Family	G4G5	S3			3	
			Species Occurrences verified in these Counties: Granite, Lincoln, Missoula, Ravalli State Rank Reason: Ligusticum verticillatum occurs in northern Idaho, western Montana, and British Columbia. It has been found in Lincoln and Ravalli Counties, growing in moist forests and meadows of spruce-fir habitats, becoming common in Idaho. Herbarium specimens from Missoula and Granite Counties may be mis-identified. Current data on locations, population sizes, and threats is greatly needed.					
Lobelia kalmii Kalm's Lobelia		Campanulaceae Bellflower Family	G5	S3			3	
			Species Occurrences verified in these Counties: Deer Lodge, Flathead, Lake, Lincoln, Powell, Sheridan, Teton, Wheatland State Rank Reason: Lobelia kalmii occurs in fens and other high-organic wetlands in northwest, central, and northeast Montana. Approximately 34 observations have been made at about 23 unique locations. The central Montana location has not been observed since 1934. Current observation, population size, and threat information at documented sites is needed.					
Lobelia spicata Pale-spiked Lobelia		Campanulaceae Bellflower Family	G5	S2?			3	Moist meadows
			Species Occurrences verified in these Counties: Dawson, Richland, Sheridan State Rank Reason: Rare and peripheral in Montana, where it is known from a few locations in the northeast corner of the state. Additional data on population levels and trends are needed. Unclear if any of the documented occurrences are subject to negative impacts or disturbances.					
Lomatium attenuatum Taper-tip Desert-parsley		Apiaceae Parsley/Carrot Family	G3	S3			SENSITIVE	3 Slopes and Scree (Dry)
			Species Occurrences verified in these Counties: Beaverhead, Madison State Rank Reason: Lomatium attenuatum is restricted to northwest Wyoming and southwest Montana, with most of its range in Montana. It is known from several locations in Beaverhead and Madison counties. Some populations may be vulnerable to impacts from mining activities and noxious weed invasion.					
Lomatium geyeri Geyer's Biscuitroot		Apiaceae Parsley/Carrot Family	G4	S2		Sensitive - Known on Forests (KOOT)	4	Rocky sites (Mesic)
			Species Occurrences verified in these Counties: Lincoln State Rank Reason: Geyer's biscuitroot occurs in northwest Montana in less than a dozen occurrences, including several large, extensive populations. Encroachment of invasive weeds from nearby infestations into habitat occupied by the species is the primary concern.					
Lomatium nuttallii Nuttall Desert-parsley		Apiaceae Parsley/Carrot Family	G3	S2			SENSITIVE	2 Rocky, pine woodlands
			Species Occurrences verified in these Counties: Big Horn, Rosebud State Rank Reason: The few populations of Nuttall's desert-parsley in the upper Tongue River drainage of Montana are disjunct from the main range of the species in southeastern Wyoming and adjacent Nebraska and Colorado. Its position on mid and lower slopes along drainages in conjunction with its occurrence on private land may make it susceptible to negative impacts from development activities. Potential future coal and/or coalbed methane development could eventually impact the species. Weeds are not currently a problem at any of the known sites. Additional locations are likely to be found in the vicinity of the known occurrences with additional surveys.					
Lomatogonium rotatum Marsh Felwort		Gentianaceae Gentians	G5	S1S2			3	Wetland/Riparian
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: Only two known occurrences in Montana on BLM and private lands, including one moderate-sized population. Livestock grazing occurs in the occupied habitat, though it is unclear what effect it may have on <i>L. rotatum</i> . Changes in the hydrology, particularly lowering of the water table may adversely affect populations.					
Malacothrix torreyi Desert Dandelion		Asteraceae Aster/Sunflowers	G4	S1S2			3	Open slopes (low-elevation)
			Species Occurrences verified in these Counties: Carbon State Rank Reason: Desert dandelion is limited in Montana to a few localized sites on the south side of the Pryor Mountains. Impacts of grazing are unknown, but it may respond positively to moderate levels of disturbance. Additional data on population levels and trends are needed.					
Mentzelia nuda Bractless blazingstar		Loasaceae Blazingstar / Stickleaf Family	G5	S1S2			3	Open areas (sandy or gravelly soils)
			Species Occurrences verified in these Counties: Big Horn, Custer, Dawson, Powder River, Roosevelt, Rosebud, Valley State Rank Reason: Rare and peripheral in Montana, where it is known from a few locations in the eastern half of the state. Additional data on population levels and trends are needed.					
Mentzelia pumila Dwarf mentzelia		Loasaceae Blazingstar / Stickleaf Family	G4	S2S3			2	Shrublands (Dry, sandy soils)
			Species Occurrences verified in these Counties: Big Horn, Carbon State Rank Reason: Rare in Montana, where it is known only from sandy sites within the Bighorn Basin area. Additional data on population levels and trends are needed.					
Mertensia bella Oregon Bluebells		Boraginaceae Borage Family	G4	S2S3		Sensitive - Known on Forests (LOLO)	3	Vernally moist soil (Montane)

			Species Occurrences verified in these Counties: Missoula State Rank Reason: Rare in Montana, where it is known only from the Lolo National Forest. Some disturbance may be beneficial or at least tolerated. Mining activity occurs near one site though it is unknown if this has had any impact on <i>M. bella</i> . Additional monitoring of the populations is needed to determine trends.						
Micranthes apetal Tiny Swamp Saxifrage	Saxifraga integrifolia Hook. var. apetal , Saxifraga apetal	Saxifragaceae Saxifrage Family	G3Q	S2?				3	Alpine
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Granite, Madison, Silver Bow State Rank Reason: Known from two occurrences, one in the East Pioneers and one in the Absaroka-Beartooth Wilderness. Both occurrences are known from single specimen collections. Though little data are available for the species in Montana, the alpine habitat in which it grows is not generally subject to negative impacts from human disturbance.						
Micranthes tempesta Storm Saxifrage	Saxifraga tempesta	Saxifragaceae Saxifrage Family	G2G3	S2S3		Sensitive - Known on Forests (BD, BRT)		3	Alpine
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Granite, Ravalli State Rank Reason: State endemic known from approximately a dozen extant sites in southwest Montana. The high elevation habitat of the species in conjunction with approximately half of the populations in designated wilderness areas minimize the potential for negative impacts to the species.						
Mimulus ampliatus Stalk-leaved Monkeyflower	Mimulus patulus , Mimulus washingtonensis	Phrymaceae Lopseed Family	G3	S3		Sensitive - Known on Forests (KOOT)		4	Vernally moist soil (Valleys to subalpine)
			Species Occurrences verified in these Counties: Beaverhead, Flathead, Glacier, Lincoln, Missoula, Park, Ravalli, Sanders State Rank Reason: See rank details.						
Mimulus breviflorus Short-flowered Monkeyflower		Phrymaceae Lopseed Family	G4	S1S2		Sensitive - Known on Forests (KOOT) Species of Conservation Concern on Forests (FLAT)		2	Rock/Talus (Mesic, Montane)
			Species Occurrences verified in these Counties: Beaverhead, Flathead, Glacier, Lincoln, Sanders State Rank Reason: Rare in Montana, where it is known from a few, scattered locations in the northwest corner of the state.						
Mimulus clivicola North Idaho Monkeyflower		Phrymaceae Lopseed Family	G4	S2?		Sensitive - Known on Forests (LOLO) Sensitive - Suspected on Forests (KOOT)		4	
			Species Occurrences verified in these Counties: Mineral, Sanders State Rank Reason: See rank details.						
Mimulus floribundus Floriferous Monkeyflower		Phrymaceae Lopseed Family	G5	SH				4	
			Species Occurrences verified in these Counties: Beaverhead, Cascade, Flathead, Glacier, Lincoln, Park, Ravalli, Sanders, Stillwater						
Mimulus hymenophyllus Thinsepal monkeyflower		Phrymaceae Lopseed Family	G2	S1S2			SENSITIVE	4	
			Species Occurrences verified in these Counties: Carbon, Lake, Park, Stillwater State Rank Reason: See rank details. Surveys of the previous collection sites are needed to document the species' status. Without additional data, a rank of "SH" will be applicable.						
Mimulus nanus Dwarf Purple Monkeyflower		Phrymaceae Lopseed Family	G5	S2S3		Sensitive - Known on Forests (BRT, CG)		1	Open slopes (low-elevation)
			Species Occurrences verified in these Counties: Gallatin, Ravalli State Rank Reason: <i>Mimulus nanus</i> is only known from a few extant occurrences in the state, plus two historical collections. Populations are generally small and in habitats susceptible to weed invasion. At least a few of the occurrences contain scattered spotted knapweed plants.						
Mimulus primuloides Primrose Monkeyflower		Phrymaceae Lopseed Family	G4	S3		Sensitive - Known on Forests (BD, BRT)		3	Fens and wet meadows
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Gallatin, Ravalli State Rank Reason: Known from several watersheds in southwest Montana, occurring almost entirely on National Forest lands. Eight of the occurrences are moderate to large-sized populations. Two historical locations are also known. Fire may adversely impact <i>M. primuloides</i> though more study is needed. It is also vulnerable to changes in hydrology and one population could be adversely affected by activity at an adjacent ski area.						
Mimulus ringens Square-stem Monkeyflower		Phrymaceae Lopseed Family	G5	S2?				3	Wetland/Riparian
			Species Occurrences verified in these Counties: Cascade, Chouteau, Fergus State Rank Reason: Rare. Currently known from a few riparian sites along the Missouri River in central Montana. Additional survey data are needed.						
Myriophyllum quitense Andean Water-milfoil		Haloragaceae Water Milfoils	G4?	S3				3	

			Species Occurrences verified in these Counties: Broadwater, Gallatin, Madison State Rank Reason: <i>Myriophyllum quitense</i> is an aquatic plant that has recently (2008-2016) been found in three waterbodies of Montana. Plants are found in slow-moving rivers that vary in water quality from the Madison River in Yellowstone National Park to Toston Reservoir on the Missouri River. These locations represent a very narrow geographical portion of Montana. Proper identification of <i>Myriophyllum</i> species require careful collections to obtain flowering or fruiting structures, use of an appropriate and current taxonomic key, and time spent studying the specimen. More surveys are greatly needed to assess the true abundance and distribution of <i>Myriophyllum quitense</i> in Montana.					
Nama densusm Nama		Hydrophyllaceae Waterleaf Family	G5	S1S2			1	Sagebrush (Sandy soil)
			Species Occurrences verified in these Counties: Carbon State Rank Reason: Nama occurs in Montana on the northeastern edge of its range. It has been found at a single location on the south side of the Pryor Mountains in 1991, occupying less than one acre of habitat. Additional survey data are needed.					
Navarretia divaricata Divaricate Navarretia		Polemoniaceae Phlox Family	G5	S1S2			3	
			Species Occurrences verified in these Counties: Sanders					
Noccaea parviflora Small-flowered Pennycress	Thlaspi parviflorum	Brassicaceae Mustards	G3	S3			3	Meadows (Moist, Montane to alpine)
			Species Occurrences verified in these Counties: Beaverhead, Carbon, Cascade, Madison, Meagher, Mineral, Park, Silver Bow State Rank Reason: <i>Noccaea parviflora</i> is a regional endemic, known in Montana from several southwestern counties. It is a small, short-lived plant that likely requires some disturbance to maintain its habitat.					
Nuttallanthus texanus Blue Toadflax	Linaria canadensis var. texana	Plantaginaceae Plantain Family	G4G5	S1S2			2	Grasslands/woodlands (sandy to clay soils)
			Species Occurrences verified in these Counties: Carter, Dawson State Rank Reason: Known from one extant occurrence in southeastern Montana near Alzada and another occurrence from Makoshika State Park. Additional surveys and monitoring are needed.					
Nymphaea leibergii Pygmy Water-lily	Nymphaea tetragona ssp. leibergii	Nymphaeaceae Water-lily Family	G5	S1			3	Aquatic
			Species Occurrences verified in these Counties: Flathead, Lake, Missoula State Rank Reason: Known from 4 extant occurrences in western valleys and one historical collection from Salmon Lake in the Seeley Lake area. Populations are susceptible to impacts from development, recreation, siltation and aquatic weeds.					
Oenothera pallida ssp. pallida Pale Evening-primrose	Oenothera pallida var. idahoensis	Onagraceae Evening-primrose Family	G5T4Q	S1			2	Sandy sites
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: Limited in Montana to the sandhills of the Centennial Valley in Beaverhead County. A reduction in natural disturbances, including fire, ungulate grazing and pocket gopher activity has led to greater dune stabilization and reduced the extent of early successional (blowout) habitat in the area.					
Oxytropis campestris var. columbiana Columbia Locoweed	Oxytropis columbiana	Fabaceae Pea Family	G5T2	S1			1	Wetland/Riparian, Gravelly shoreline
			Species Occurrences verified in these Counties: Lake State Rank Reason: Originally known in Montana from six occurrences all around Flathead Lake. However, two of the occurrences are now extirpated. Private lands, which are subject to development in the area, play a vital role in maintaining viable populations of this plant in Montana.					
Oxytropis deflexa var. foliolosa Nodding Locoweed		Fabaceae Pea Family	G5T5	S2S3			3	Alpine
			Species Occurrences verified in these Counties: Beaverhead, Gallatin, Madison, Park State Rank Reason: Rare in Montana, where it has been documented from a few, high-elevation sites in the mountains of the southwest portion of the state.					
Oxytropis parryi Parry's Locoweed		Fabaceae Pea Family	G5	S2S3			3	Alpine
			Species Occurrences verified in these Counties: Beaverhead, Madison State Rank Reason: Rare in Montana where it is known only from a few occurrences in the southwestern portion of the state. However, the species high-elevation habitat and its viability do not appear to be at significant risk at the current time.					
Oxytropis podocarpa Stalked-pod Locoweed		Fabaceae Pea Family	G4G5	S1		Sensitive - Suspected on Forests (BD)	3	Alpine
			Species Occurrences verified in these Counties: Glacier, Teton State Rank Reason: Rare in Montana, where it is known from a small area of the Rocky Mountain Front. The remote habitat should limit the possibility of negative impacts.					
Papaver pygmaeum Alpine Glacier Poppy	Papaver radicatum var. pygmaeum	Papaveraceae Poppy Family	G3	S2S3			3	Alpine
			Species Occurrences verified in these Counties: Flathead, Glacier, Lewis and Clark State Rank Reason: See rank details.					
Papaver radicatum ssp. kluanensis Alpine Poppy	Papaver kluanense, Papaver kluenensis	Papaveraceae Poppy Family	G5T4	S2S3			3	Alpine
			Species Occurrences verified in these Counties: Carbon, Park, Sweet Grass State Rank Reason: See rank details.					

Pedicularis contorta var. ctenophora Pink Coil-beaked Lousewort		Orobanchaceae Broomrape Family	G5T3	S2S3				3	Slopes (Montane/Subalpine)
			Species Occurrences verified in these Counties: Beaverhead, Judith Basin, Madison, Ravalli, Teton State Rank Reason: Restricted to extreme southwestern Montana where it is documented from a few populations. Limited data is available for the species and it may be more common than the few collections indicate.						
Pedicularis contorta var. rubicunda Selway Coil-beaked Lousewort		Orobanchaceae Broomrape Family	G5T3	S2S3				3	Ridgetops and meadows (subalpine and alpine)
			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Restricted in Montana to the Bitterroot Mountains where it is documented from several occurrences. Limited data are available for the species and it may be more common than the few collections indicate.						
Pedicularis crenulata Scallop-leaf Lousewort		Orobanchaceae Broomrape Family	G4	S1			SENSITIVE	1	Wetland/Riparian
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: Two known populations in Montana. Much of the riparian meadow habitat occupied by this species has been converted to agriculture or is being used as hay meadows.						
Pedicularis pulchella Mountain Lousewort		Orobanchaceae Broomrape Family	G3	S3				3	Alpine
			Species Occurrences verified in these Counties: Carbon, Deer Lodge, Gallatin, Granite, Madison, Park, Powell, Sweet Grass State Rank Reason: Restricted to high elevation areas of southern Montana. Limited data are available for the species and it may be more common than the few collections indicate.						
Penstemon angustifolius Narrowleaf Penstemon		Plantaginaceae Plantain Family	G5	S2S3				3	Sandy sites
			Species Occurrences verified in these Counties: Carter, Dawson, Fallon, Granite State Rank Reason: Over a dozen, small extant and/or presumed extant occurrences are known in southeast Montana, plus a few historical collections from the same area. Only one of the known populations appears to be relatively large. Additional suitable, but unsurveyed habitat likely exists in eastern Montana.						
Penstemon caryi Cary's Beardtongue		Plantaginaceae Plantain Family	G3	S3			SENSITIVE	3	Grasslands and slopes (Open, montane)
			Species Occurrences verified in these Counties: Carbon State Rank Reason: Restricted in Montana to the Pryor Mountains.						
Penstemon flavescens Yellow Beardtongue		Plantaginaceae Plantain Family	G3	S3				3	Rocky slopes (Open, montane)
			Species Occurrences verified in these Counties: Mineral, Missoula, Ravalli State Rank Reason: Restricted in Montana to the Bitterroot Range primarily in Ravalli County but also documented from Mineral County. The species can be relatively common or widely scattered in areas of suitable habitat, though detailed information on the abundance of the species is lacking. More detailed information documenting the abundance, distribution and any potential threats is needed.						
Penstemon grandiflorus Large Flowered Beardtongue		Plantaginaceae Plantain Family	G5?	S1				4	Sandy soils
			Species Occurrences verified in these Counties: Custer State Rank Reason: Rare in Montana, where it is known from only a few sites on the plains of eastern Montana.						
Penstemon humilis Low Beardtongue		Plantaginaceae Plantain Family	G5	S1S3				4	Sagebrush steppe (Montane)
			Species Occurrences verified in these Counties: Beaverhead, Gallatin, Lewis and Clark, Lincoln, Madison, Meagher, Missoula, Park, Powell, Ravalli State Rank Reason: Known in Montana from 1 collection from Beaverhead County						
Penstemon lemhiensis Lemhi Beardtongue		Plantaginaceae Plantain Family	G3	S3		Sensitive - Known on Forests (BD, BRT)	SENSITIVE	2	Sagebrush-grasslands
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Ravalli, Silver Bow State Rank Reason: <i>Penstemon lemhiensis</i> is a regional endemic that occurs only in southwest Montana and adjacent Idaho. There are numerous occurrences in Beaverhead and Ravalli Counties with a few additional occurrences located in Deer Lodge and Silver Bow Counties in Montana, but most are small to moderate in size. The number of plants in Montana is estimated at approximately 10,000 individual plants based on recent survey efforts. Plants occur on a mix of federal, state and private ownerships with National Forest lands supporting the majority of the occurrences. The species is primarily sensitive to negative impacts associated with drought conditions and fire suppression, both of which are believed to have played a significant role in the species' decline. Additional impacts to populations are occurring from noxious weed invasion, primarily spotted knapweed in the Bitterroot region. Heavy livestock grazing also negatively impacts the species. Several occurrences are found adjacent to roadsides and thus may be impacted by activities associated with road construction, maintenance and use.						
Penstemon payettensis Payette Beardtongue		Plantaginaceae Plantain Family	G4	S1		Sensitive - Known on Forests (BRT)		1	Slopes (Open, Montane)
			Species Occurrences verified in these Counties: Beaverhead, Ravalli State Rank Reason: Known in Montana from only two small occurrences in close proximity on the Bitterroot National Forest. Spotted knapweed invasion, fire suppression and road construction/maintenance are all concerns for the viability of the species in Montana. Additional data on the species in Montana are needed.						
Penstemon whippleanus Whipple's Beardtongue		Plantaginaceae Plantain Family	G5	S2				3	Open areas (subalpine and alpine)

			Species Occurrences verified in these Counties: Beaverhead, Gallatin, Madison State Rank Reason: Whipple's beardtongue occurs at the edge of its range in Montana, and is known here from just two collections, only one of which is recent. The species occupies high elevation, rocky habitat that is relatively unthreatened.					
Petasites frigidus var. frigidus Arctic Sweet Coltsfoot		Asteraceae Aster/Sunflowers	G5T5	S2		Species of Conservation Concern on Forests (FLAT)	2	Wetland/Riparian
			Species Occurrences verified in these Counties: Flathead, Glacier State Rank Reason: Rare in Montana, where it is at the southern edge of its range. Known from a few widely scattered sites in the northwest corner of the state.					
Phacelia incana Hoary Phacelia		Hydrophyllaceae Waterleaf Family	G3G4	S3			3	Rocky slopes (foothills)
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: <i>Phacelia incana</i> occurs in Idaho, Nevada, Utah, Colorado and Montana. In Montana, it is known from approximately ten occurrences in Beaverhead County. It is difficult to estimate the size of populations because the plant is an annual, and seed germination varies greatly with climate. Habitat is probably not threatened by anthropogenic sources.					
Phacelia thermalis Hot Spring Phacelia		Hydrophyllaceae Waterleaf Family	G3G4	S1S3			4	Barren clay slopes
			Species Occurrences verified in these Counties: Fergus, Garfield, Phillips, Valley State Rank Reason: Hot spring phacelia is known from a very small number of sites in northeastern Montana, where it is disjunct from its primary range (northern California to southwestern Idaho). The species is an annual and may be vulnerable to competition from invasive exotics, particularly sweet clover, which is widespread in the type of habitat where hot spring phacelia has been found.					
Phlox kelseyi var. missoulensis Missoula Phlox	Phlox missoulensis	Polemoniaceae Phlox Family	G3	S3		Sensitive - Known on Forests (BD) Sensitive - Suspected on Forests (LOLO) Species of Conservation Concern on Forests (HLC)	1	Slopes/ridges (Open, foothills to subalpine)
			Species Occurrences verified in these Counties: Cascade, Granite, Jefferson, Judith Basin, Lewis and Clark, Madison, Meagher, Missoula, Powell, Teton State Rank Reason: Missoula phlox is a state endemic known from over 2 dozen occurrences in west-central Montana, most of which are moderate to large-sized. Populations occur on a mix of ownerships, including private lands which host several occurrences. The Waterworks Hill population is infested with several noxious weeds and heavy recreational trail use also occurs within the occupied habitat. Other populations appear to be at much less risk though some impacts from invasive weeds, recreational use and development are possible.					
Physaria brassicoides Double Bladderpod		Brassicaceae Mustards	G5	S3			3	Breaklands/badlands
			Species Occurrences verified in these Counties: Carbon, Carter, Custer, Petroleum, Phillips, Powder River, Stillwater State Rank Reason: Double bladderpod is endemic to a restricted area of the northern Great Plains, and is known in Montana only from a handful of populations. Populations occur on a mix of federal, state and private ownerships. Impacts to the species from livestock grazing and invasive weeds are minimal at this time as the typically steep, sparsely-vegetated habitat is not conducive to grazing. Yellow sweetclover was observed at one location and it may eventually have a negative impact on the species.					
Physaria carinata Keel-bladderpod	Lesquerella carinata, Lesquerella carinata var. languida, Lesquerella paysonii [misapplied in MT], Physaria carinata ssp. carinata	Brassicaceae Mustards	G3G4TNR	S1S2		Sensitive - Known on Forests (BD)	1	Grassland slopes (low-elevation)
			Species Occurrences verified in these Counties: Beaverhead, Granite State Rank Reason: <i>Physaria carinata</i> is restricted to areas of calcareous limestone substrates on low elevation, south-facing grasslands of Granite and Beaverhead Counties. Population numbers appear to have declined significantly in at least several of the occurrences in the Garnet Mountains from the time they were first documented in the 1980's and early 1990's. During this time period, spotted knapweed densities have increased in the area and the noxious weed is now a dominant plant in most of the keel-bladderpod sites. At least one previous study has documented decreased vigor and survivorship of keel-bladderpod in knapweed infested areas.					
Physaria didymocarpa var. lanata Woolly Twinpod		Brassicaceae Mustards	G5T2	S2S3			4	Grasslands/Shrublands (Open, plains)
			Species Occurrences verified in these Counties: Big Horn, Rosebud State Rank Reason: Only a few known occurrences in Montana, including two potentially large populations. However, lots of unsurveyed potential habitat exists. Both BLM and private lands are important to the viability of the species in Montana. Oil and gas development, coalbed methane, and invasive weeds have the potential to detrimentally impact populations.					
Physaria douglasii Douglas Bladderpod	Lesquerella douglasii	Brassicaceae Mustards	G5	S1			1	Woodlands (Sandy soils, low-elevation)
			Species Occurrences verified in these Counties: Lincoln State Rank Reason: Known from one population in northwest Montana at the edge of Lake Koocanusa. Impacts to the population from ORV use, recreation and erosion of the sandy bluffs are possible, though additional monitoring is needed to determine what impacts if any are occurring.					

Potentilla hyparctica Low Arctic Cinquefoil	Potentilla nana, Potentilla flabellifolia var. emarginata	Rosaceae Rose Family	G5	S2			3	Alpine
Species Occurrences verified in these Counties: Carbon State Rank Reason: Rare in Montana, where it is currently only from a couple collections from the Beartooth Mtns. The remote, high-elevation habitat should greatly minimize the potential for any negative impacts to the viability of the species in the state. Accurate estimates of population levels are lacking.								
Potentilla nivea var. pentaphylla Five-leaf Cinquefoil	Potentilla quinquefolia	Rosaceae Rose Family	G5T4	S3		Sensitive - Known on Forests (BD)	4	Alpine
Species Occurrences verified in these Counties: Flathead, Glacier, Lincoln, Madison, Park, Pondera State Rank Reason: Rare in Montana, though several large populations are known and most populations, as well as the species' habitat, are not being negatively impacted.								
Potentilla plattensis Platte Cinquefoil		Rosaceae Rose Family	G4	S3			4	Grasslands/Sagebrush (Mesic)
Species Occurrences verified in these Counties: Beaverhead, Carbon, Judith Basin, Valley State Rank Reason: Rare in Montana, where it is known from several collections, particularly from Beaverhead County.								
Primula alcalina Alkali Primrose		Primulaceae Primrose Family	G2	S2		Sensitive - Known on Forests (BD)	1	Wetland/Riparian
Species Occurrences verified in these Counties: Beaverhead, Madison State Rank Reason: <i>Primula alcalina</i> is a regional endemic, occurring only in east-central Idaho and adjacent Montana, where it is known from just one recently documented population in Beaverhead County on BLM and National Forest lands. Another population documented by a historical collection from 1920 by F. Rose has not been relocated. The extant location is actively grazed and the species may be vulnerable to impacts associated with cattle grazing and activities that alter the hydrology (irrigation, diversions).								
Primula incana Mealy Primrose		Primulaceae Primrose Family	G5	S3		Sensitive - Known on Forests (BD) Sensitive - Historically known, not recently documented on Forests (CG)	2	Wetland/Riparian
Species Occurrences verified in these Counties: Beaverhead, Broadwater, Carbon, Deer Lodge, Gallatin, Jefferson, Madison, Meagher, Powell, Sheridan, Silver Bow, Teton State Rank Reason: <i>Primula incana</i> is known from a few dozen extant occurrences in Montana, including several moderate to large populations. However, most known populations are small, and the status of several populations is uncertain. Ownership of the occupied areas is varied and includes federal, state and private lands, including several locations managed or protected for their conservation values. However, unprotected private lands host many occurrences. Cattle grazing may have some negative effects on the species including the direct effects of herbivory and trampling. The species is also vulnerable to activities that alter the hydrology of the wetlands it occupies. Continued threats and potentially declining trends, particularly in regards to habitat quality make the species' vulnerable to local extirpation.								
Prunus pumila Sand Cherry		Rosaceae Rose Family	G5	S1S3			2	Sandy or rocky soils (Plains)
Species Occurrences verified in these Counties: Fallon State Rank Reason: The sole known extant location in Montana occurs along a county road and is susceptible to road construction and maintenance activities. A 1960 collection with vague locational data has not been relocated but it apparently occurred in native habitat.								
Psilocarphus brevissimus Dwarf woolly-heads		Asteraceae Aster/Sunflowers	G4	S2S3		Sensitive - Known on Forests (KOOT)	3	Wetland/Riparian
Species Occurrences verified in these Counties: Cascade, Lincoln, Petroleum, Phillips, Sanders, Valley State Rank Reason: Limited data combined with the possibility that several reported observations from western MT may be mis-identified with other Psilocarphus species make a precise determination of the species' status difficult.								
Pyrrocoma carthamoides var. subsquarrosa Beartooth Large-flowered Goldenweed	Haplopappus carthamoides var. subsquarrosus	Asteraceae Aster/Sunflowers	G4G5T3	S3		Sensitive - Known on Forests (CG)	3	Sagebrush-Grassland
Species Occurrences verified in these Counties: Carbon State Rank Reason: The Beartooth large-flowered goldenweed is a local endemic to the eastern front of the Beartooth Mountains and the foothills of the Pryor Mountains and adjacent areas of Wyoming. Although several populations are large, it is vulnerable to increased shrub and tree cover due to fire suppression and to competition from invasive plants.								
Quercus macrocarpa Bur Oak		Fagaceae Beech / Oaks	G5	S2			2	Shale ridges
Species Occurrences verified in these Counties: Carter State Rank Reason: Bur oak is at the extreme western edge of its range in Montana, where it occurs in a localized, though fairly large, occurrence in Carter County. Bentonite mining is active in this area and exotic weeds are prevalent though negative impacts to bur oak have not been documented due to a lack of surveys and monitoring.								
Ranunculus cardiophyllus Heart-leaved Buttercup		Ranunculaceae Buttercup Family	G5	S3			3	Grasslands (Moist, Montane)
Species Occurrences verified in these Counties: Chouteau, Glacier, Sweet Grass, Toole State Rank Reason: Rare in Montana, where it is primarily distributed in the north-central part of the state.								

Ranunculus grayi Arctic Buttercup	Ranunculus karelinii, Ranunculus verecundus, Ranunculus gelidus	Ranunculaceae Buttercup Family	G4G5	S3				3	Alpine
			Species Occurrences verified in these Counties: Carbon, Deer Lodge, Flathead, Glacier, Madison, Meagher, Park, Stillwater State Rank Reason: Also includes <i>R. verecundus</i> , which was formerly tracked as a separate Species of Concern.						
Ranunculus orthorhynchus Straightbeak Buttercup		Ranunculaceae Buttercup Family	G5	S1S2				2	Wetland/Riparian (Montane)
			Species Occurrences verified in these Counties: Deer Lodge, Flathead, Glacier, Granite, Lake, Mineral, Missoula, Sanders State Rank Reason: Rare in Montana, where it is known from the western portion of the state based upon several specimen collections. However, only one collection has been made in the past two decades. Additional data are needed to determine this species' status.						
Ranunculus pedatifidus Northern Buttercup		Ranunculaceae Buttercup Family	G5	S3		Species of Conservation Concern on Forests (HLC)		3	Meadows/Woodlands (Montane to Alpine)
			Species Occurrences verified in these Counties: Carbon, Flathead, Glacier, Liberty, Sweet Grass, Teton State Rank Reason: Rare in Montana. Documented in the state from several collections. Additional data are needed to more precisely determine the species' status.						
Ribes laxiflorum Trailing Black Currant		Grossulariaceae Currants / Gooseberries	G5	S2?				4	Shrublands (Rocky, montane)
			Species Occurrences verified in these Counties: Lincoln State Rank Reason: Rare in Montana, where it is known from a single collection from Lincoln County. The documented population does not appear to be at risk. Additional data are needed.						
Ribes triste Swamp Red Currant		Grossulariaceae Currants / Gooseberries	G5	S2?				4	Forest openings (Mesic, montane/subalpine)
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Granite, Mineral, Ravalli State Rank Reason: Rare in Montana, where it is known from a few collections from the western portion of the state. Additional data are needed.						
Rorippa calycina Persistent-sepal Yellow-cress		Brassicaceae Mustards	G3	SH				4	Wetland/Riparian
			Species Occurrences verified in these Counties: Big Horn, Custer, Mccone, Rosebud, Treasure, Yellowstone State Rank Reason: <i>Rorippa calycina</i> is a regional endemic currently known only from four Montana records. The species was last observed in Montana more than 30 years ago. Surveys are needed.						
Rotala ramosior Toothcup		Lythraceae Loosestrife Family	G5	S1S2				4	Wetland/Riparian
			Species Occurrences verified in these Counties: Lake, Missoula, Ravalli State Rank Reason: Rare in Montana, where it is known from approximately a half-dozen wetland sites in the valley bottoms in the western portion of the state. Potential threats and impacts to the known occurrences, as well as population trends, need to be evaluated.						
Rubus arcticus Nagoonberry	Rubus acaulis, Rubus arcticus ssp. acaulis	Rosaceae Rose Family	G5	S2				3	
			Species Occurrences verified in these Counties: Beaverhead, Flathead, Glacier, Lincoln, Madison State Rank Reason: <i>Rubus acaulis</i> may be rare or common where its habitat is present. However, its habitat (hummocks in <i>Sphagnum</i> -moss dominated fens, high elevation wet-meadows, etc.) is very specific and often limited in Montana.						
Sagina nivalis Arctic Pearlwort		Caryophyllaceae Pink Family	G5	S2S3				3	Alpine
			Species Occurrences verified in these Counties: Carbon, Glacier, Stillwater State Rank Reason: Rare in Montana, where it is known from Glacier National Park and the Beartooth Plateau. The remote, high-elevation habitat should greatly minimize the potential for any negative impacts to the viability of the species in the state. Accurate estimates of population levels are lacking.						
Salix barrattiana Barratt's Willow		Salicaceae Willows / Poplar	G5	S2		Sensitive - Known on Forests (CG)		3	Alpine
			Species Occurrences verified in these Counties: Carbon, Glacier, Madison State Rank Reason: Rare in Montana. Known from two disjunct sites, one in Glacier National Park and one on the Beartooth Plateau. Populations are small, but the remote, high-elevation habitat should greatly minimize the potential for any negative impacts to the viability of the species in the state.						
Salix cascadenis Cascade Willow		Salicaceae Willows / Poplar	G5	S2				3	Alpine
			Species Occurrences verified in these Counties: Deer Lodge State Rank Reason: Rare in Montana. Species is known in Montana only from a small area of the Anaconda-Pintlers. The remote, high-elevation habitat should greatly minimize the potential for any negative impacts to the viability of the species in the state. Accurate estimates of population levels are lacking.						
Salix serissima Autumn Willow		Salicaceae Willows / Poplar	G5	S3				3	Wetland/Riparian
			Species Occurrences verified in these Counties: Cascade, Glacier, Meagher, Pondera, Teton State Rank Reason: This willow is primarily found in Montana along the Rocky Mountain Front. Approximately half the occurrences are on lands managed in part for their conservation value. The species is primarily susceptible to impacts associated with heavy grazing and changes in the hydrology of the fens and wet meadows which it occupies.						
Sandbergia perplexa Puzzling Rockcress	Halimolobos perplexa	Brassicaceae Mustards	G4	S2		Sensitive - Known on Forests (BRT)		2	Shrubland/woodland slopes (Open, Montane)

			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Rare in Montana, where it is known only from the very southern end of the Bitterroot Valley on the Bitterroot National Forest. Spotted knapweed is known from at least one of the populations and further spread of invasive weeds at the known occurrences is likely without control measures. Trend data and repeat observations of the known occurrences are lacking.					
Satureja douglasii Yerba Buena	Clinopodium douglasii	Lamiaceae Mints	G5	S3			3	Forest (Moist, montane)
			Species Occurrences verified in these Counties: Mineral, Missoula, Ravalli, Sanders State Rank Reason: Rare in Montana, where it is known from several sites near the Idaho border. It is primarily a coastal species, disjunct in western Montana. Population levels appear healthy and may be increasing in some areas.					
Saussurea nuda Dwarf Saw-wort	Saussurea nuda var. densa, Saussurea densa	Asteraceae Aster/Sunflowers	G4	S2S3			3	Alpine
			Species Occurrences verified in these Counties: Flathead, Lewis and Clark, Pondera, Teton State Rank Reason: Known from one handfull of small occurrences along the Rocky Mountain Front, primarily in the Bob Marshall Wilderness Complex. Limited data are available for most occurrences leading to the uncertainty in the species' rank.					
Saussurea weberi Weber's Saw-wort		Asteraceae Aster/Sunflowers	G3	S2		Sensitive - Known on Forests (BD)	3	Alpine
			Species Occurrences verified in these Counties: Deer Lodge, Granite, Park State Rank Reason: Known from one large occurrence in the Anaconda-Pintler Range in the alpine zone. The remote, high-elevation habitat should greatly minimize the potential for any negative impacts to the viability of the species in the state. Population estimates from the single, documented occurrence vary widely. Additional population data are needed.					
Saxifraga hirculus Yellow Marsh Saxifrage		Saxifragaceae Saxifrage Family	G5	S1S2			3	Alpine
			Species Occurrences verified in these Counties: Carbon State Rank Reason: Known from one small population in the Absorka-Beartooth Wilderness. Though little data are available for the species in Montana, the alpine habitat in which it grows is not generally subject to negative impacts from human disturbance.					
Senecio amplexens Clasping Groundsel	Ligularia amplexens	Asteraceae Aster/Sunflowers	G4	S1S2			4	Alpine
			Species Occurrences verified in these Counties: Carbon, Glacier State Rank Reason: In Montana, only known from the Beartooth (Line Creek) Plateau. Additional data on population size, trends and potential threats are needed to evaluate the species' vulnerability.					
Senecio elmeri Elmer's Ragwort	Senecio sribillei	Asteraceae Aster/Sunflowers	G4	S2			3	Alpine
			Species Occurrences verified in these Counties: Lincoln, Sanders State Rank Reason: Rare in the state. Known from only one high-elevation site in the Cabinet Mountains. Its location in a designated wilderness and its high-elevation habitat should prevent most detrimental impacts to the species' viability in Montana.					
Senecio eremophilus Desert Groundsel		Asteraceae Aster/Sunflowers	G5	S1S2			3	Wetland/Riparian
			Species Occurrences verified in these Counties: Big Horn, Blaine, Flathead, Glacier, Hill, Lake, Lincoln, Phillips State Rank Reason: Known from at least 5 occurrences, including two historical collections. Little data are available for this species in Montana. More information is needed. May be more common than collections indicate.					
Senecio hydrophilus Alkali-marsh Ragwort		Asteraceae Aster/Sunflowers	G5	S3			3	
			Species Occurrences verified in these Counties: Beaverhead, Broadwater, Gallatin, Madison State Rank Reason: Senecio hydrophilus is present in alkaline habitats within a portion of southwest Montana. Plants are not that common, and occur in low-elevation wetlands that can be victim to dewatering.					
Senecio integerrimus var. scribneri Scribner's Ragwort		Asteraceae Aster/Sunflowers	G5T2T3	S2S3			3	
			Species Occurrences verified in these Counties: Carbon, Custer, Fergus, Golden Valley, Hill, Liberty, Musselshell, Park, Phillips, Rosebud, Valley, Wheatland, Yellowstone State Rank Reason: See rank details.					
Shoshonea pulvinata Shoshonea		Apiaceae Parsley/Carrot Family	G3	S2		Sensitive - Known on Forests (CG)	3	Rock Outcrops
			Species Occurrences verified in these Counties: Carbon State Rank Reason: Known in Montana only from the Pryor Mountains and the eastern slope of the Beartooth Plateau. Occurrences are located mostly on federal lands.					
Sidalcea oregana Oregon Checker-mallow	Sidalcea organa ssp. oregana	Malvaceae Mallow Family	G5	S2S3			1	Grasslands (low-elevation)
			Species Occurrences verified in these Counties: Gallatin, Lake State Rank Reason: Known from two widely separate sites in Gallatin and Lake counties. Habitats occupied by the species are susceptible to weed invasion and both locations have a large component of weedy species. However, <i>S. oregana</i> appears capable of tolerating at least some competition from these weedy species. The Lake County population occurs near and along Highway 93 and has the potential to be significantly negatively impacted by highway construction.					
Silene spaldingii Spalding's Catchfly	Spalding's Campion	Caryophyllaceae Pink Family	G2	S2	LT		1	Grasslands (Intermountain)

			Species Occurrences verified in these Counties: Flathead, Lake, Lincoln, Sanders State Rank Reason: <i>Silene spaldingii</i> exists in only a few locations in the northwest corner of the state. Extant occurrences are known in the following areas: Tobacco Plains area, Lost Trail National Wildlife Refuge, the Niarada area and on Wild Horse Island. The majority of occurrences have less than 100 individuals, though 3 sites are each known to contain over 1,000 individuals and the total population size in Montana is likely 20,000+ mature plants based upon 2011 data. One historical occurrence exists from the Columbia Falls area. Several threats affect the long-term viability of the species in the state. Invasive weeds are the most widespread threat and are negatively impacting the bunchgrass habitat occupied by <i>S. spaldingii</i> . Housing development and subdivision are directly impacting populations in the Tobacco Plains and has the potential to further isolate known occurrences in the area. Cattle grazing is affecting several populations and two other occurrences have apparently been extirpated recently from the severe impacts associated with llama grazing. Fire exclusion and the successive build-up of litter compared to historical conditions appears to be having negative impacts on survival and reproduction. Populations are also at risk due to the small numbers of individuals and their isolated nature, which reduces the chances of cross-pollination and gene flow between populations. Long- and short-term trends are difficult to gauge due to the lack of survey and monitoring data. Estimates of trends and population size are also compounded by <i>S. spaldingii</i> plants exhibiting summer dormancy at rates that vary widely from year to year.						
Solidago ptarmicoides Prairie Goldenrod	Oligoneuron album, Aster ptarmicoides	Asteraceae Aster/Sunflowers	G5	S2S3				3	Grasslands (Plains)
			Species Occurrences verified in these Counties: Carter, Richland, Wibaux State Rank Reason: Rare in Montana, where it has been documented from only a few locations on the eastern plains.						
Sphaeromeria argentea Chicken-sage	Tanacetum nuttallii, Artemisia macarthurii	Asteraceae Aster/Sunflowers	G3G4	S3			SENSITIVE	3	Sagebrush steppe (low-elevation)
			Species Occurrences verified in these Counties: Beaverhead State Rank Reason: <i>Sphaeromeria argentea</i> occurs in east-central Idaho and adjacent Beaverhead County, Montana with disjunct populations in Nevada as well as southwest Wyoming and adjacent Colorado. There are nearly 20 known locations south of Dillon; many populations are sparse but spread over large areas, so population estimates are difficult. All known populations are subject to livestock grazing; however chicken sage is aromatic and most likely unpalatable to cattle.						
Stellaria crassifolia Fleshy Stitchwort		Caryophyllaceae Pink Family	G5	S2				3	Wetland/Riparian
			Species Occurrences verified in these Counties: Beaverhead, Glacier, Granite State Rank Reason: Rare in Montana where it is known from a few sparsely distributed locations that are mostly poorly documented.						
Sullivantia hapemanii Wyoming Sullivantia		Saxifragaceae Saxifrage Family	G3	S2S3				3	Rock/Talus
			Species Occurrences verified in these Counties: Big Horn, Carbon State Rank Reason: Wyoming Sullivantia is regional endemic known in Montana only from a few, clustered locations. It grows in small, fragile aquatic habitats that may be vulnerable to hydrologic changes from water development or diversion, or trampling.						
Symphyotrichum molle Soft Aster		Asteraceae Aster/Sunflowers	G3	S1S3				4	NA
			Species Occurrences verified in these Counties: Big Horn, Carbon State Rank Reason: Known in Montana from 1 collection from the Bighorn Mtns. Though its exact status is uncertain, its rarity warrants its inclusion as a Species of Concern.						
Synthyris canbyi Mission Mountain kittentails	Veronica canbyi	Plantaginaceae Plantain Family	G2G3	S2S3				3	Alpine
			Species Occurrences verified in these Counties: Flathead, Granite, Lake, Missoula, Ravalli State Rank Reason: State endemic with 10 occurrences restricted to high elevation, open, rocky slopes in the Mission and Swan Ranges. As such, habitat is not generally prone to human disturbance and most occurrences are in designated wilderness areas. Additional occurrences likely exist across the known range of the species.						
Tetradymia spinosa Short-spine Horsebrush		Asteraceae Aster/Sunflowers	G5	S2S3				4	
			Species Occurrences verified in these Counties: Beaverhead, Big Horn, Carbon, Park State Rank Reason: <i>Tetradymia spinosa</i> occurs in Montana on the northern edge of its distribution. Since 1900 <i>Tetradymia spinosa</i> has been included in the major floras for Montana (Rydberg 1900, Booth 1966, Dorn 1984, Lesica et al. 2012); yet, herbarium specimens are relatively few and the vast majority of observation data is more than 25 years old. In Beaverhead County its presence is documented by a single 1888 herbarium specimen and a 2013 observation with few informational details. Based on available information, this shrub grows in a common habitat type (sagebrush steppe) where populations are often small and widely scattered. In general, seedlings of <i>Tetradymia</i> species have been observed as rare, partially due to durations where flowers are not produced, germination is poor, and establishment rates are low (Strothers 1974). Surveys that bring forth better mapping and information on locations, population sizes, viability, reproduction, habitat, threats, and ecology are greatly needed.						
Thalictrum alpinum Alpine Meadowrue		Ranunculaceae Buttercup Family	G5	S2		Sensitive - Known on Forests (BD) Sensitive - Suspected on Forests (CG)		2	Wetland/Riparian
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Granite, Madison State Rank Reason: Rare in Montana, where it is known from approximately two dozen sites mostly on public land. Its habitat is vulnerable to hydrological alteration. Grazing can be beneficial, except where it leads to stream downcutting and loss of riparian habitat.						

Thelypodium paniculatum Northwestern Thelypody	Thelypodium sagittatum var. <i>crassicaarpum</i>	Brassicaceae Mustards	G2	SH			3	Wetland/Riparian
			Species Occurrences verified in these Counties: Beaverhead, Gallatin, Madison State Rank Reason: Known only from an 1899 collection in Beaverhead County, although Dorn (1984) also reports it for Madison County.					
Thelypodium sagittatum Slender Thelypody		Brassicaceae Mustards	G4	S2			3	Alkaline meadows (Valleys and Montane)
			Species Occurrences verified in these Counties: Beaverhead, Gallatin, Madison State Rank Reason: Known from numerous occurrences in extreme southwestern Montana.					
Tonestus aberrans Idaho Goldenweed	Haplopappus aberrans, Triniteurybia aberrans, Eurybia aberrans	Asteraceae Aster/Sunflowers	G3	S1S2		Sensitive - Known on Forests (BRT)	2	Rock/Talus
			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Known from two moderate-sized occurrences and two smaller occurrences on the Bitterroot National Forest and adjacent private land. One population occurs adjacent to a road, where construction may have impacted the population. No negative impacts to the populations are currently known to be occurring. However, populations are susceptible to potential impacts associated with roads and rock climbing.					
Townsendia condensata Cushion Townsend-daisy		Asteraceae Aster/Sunflowers	G4	S1S3			3	Alpine
			Species Occurrences verified in these Counties: Beaverhead, Flathead, Glacier, Park State Rank Reason: Cushion townsendia is known in Montana from one presumed extant occurrence in Glacier National Park and three other historical collections from GNP and the Beartooth Mountains. Risks are likely minimal given the remoteness of its alpine habitat.					
Townsendia florifer Showy Townsend-daisy	Townsendia florifera	Asteraceae Aster/Sunflowers	G5	S2			3	Grasslands and Sagebrush
			Species Occurrences verified in these Counties: Beaverhead, Park, Sweet Grass State Rank Reason: Known in Montana from only a few, small occurrences in the southwestern corner of the state.					
Trifolium cyathiferum Cup Clover		Fabaceae Pea Family	G4	S3			4	
			Species Occurrences verified in these Counties: Missoula, Ravalli State Rank Reason: Trifolium cyathiferum occurs in two counties with limited information on population size. One occurrence was re-visited in 1998 and found to be absent due to habitat succession.					
Trifolium eriocephalum Woolly-head Clover		Fabaceae Pea Family	G5	S2		Sensitive - Known on Forests (BRT) Sensitive - Suspected on Forests (BD, LOLO)	2	Open areas (foothills and montane)
			Species Occurrences verified in these Counties: Beaverhead, Ravalli State Rank Reason: Known from eight large occurrences on the Bitterroot National Forest. Invasive weeds, particularly spotted knapweed, are a problem in the habitat occupied by the species. Timber harvest and related road-building activities may also negatively impact populations. However, <i>Trifolium eriocephalum</i> appears capable of tolerating some level of disturbance.					
Trifolium gymnocarpon Hollyleaf Clover		Fabaceae Pea Family	G5	S2		Sensitive - Known on Forests (BRT, LOLO) Sensitive - Suspected on Forests (BD)	2	Open areas (foothills and montane)
			Species Occurrences verified in these Counties: Granite, Ravalli State Rank Reason: Known from many sites within the West Fork Bitterroot River drainage, which would encompass one large metapopulation. Also known in Montana from one disjunct occurrence in the Rock Creek drainage on the Lolo National Forest. Invasive weeds, particularly spotted knapweed, are a problem in some of the habitat occupied by the species. However, <i>Trifolium gymnocarpon</i> , as with other clover species, appears capable of tolerating or even benefitting from some disturbance.					
Trifolium microcephalum Woolly Clover		Fabaceae Pea Family	G5	S3			2	
			Species Occurrences verified in these Counties: Missoula, Ravalli					
Triodanis leptocarpa Slim-pod Venus'-looking-glass	Specularia leptocarpa	Campanulaceae Bellflower Family	G5?	S3			3	
			Species Occurrences verified in these Counties: Big Horn, Carter, Cascade, Chouteau, Custer, Park, Petroleum, Phillips, Powder River, Rosebud, Stillwater, Sweet Grass, Valley State Rank Reason: Triodanis leptocarpa is common in the southern Great Plains and extends into eastern and central Montana. It occurs in grasslands, grass-dominated rocky slopes, and sagebrush-dominated grasslands. It has been found in grazed and ungrazed lands and appears to tolerate some disturbance. Approximately 14 locations were documented prior to 1958 and occur in central Montana. Approximately 14 locations were documented since 1974 and mostly occur in eastern Montana. Re-visits to known locations and current population data is greatly needed.					
Utricularia intermedia Flatleaf Bladderwort		Lentibulariaceae Bladderworts	G5	S2		Sensitive - Known on Forests (KOOT)	3	Fens (Aquatic)
			Species Occurrences verified in these Counties: Flathead, Glacier, Lake, Lincoln, Madison, Missoula, Powell State Rank Reason: Only known from a few occurrences in the western half of the state.					

Carex chordorrhiza Creeping Sedge		Cyperaceae Sedges	G5	S3		Sensitive - Known on Forests (KOOT) Sensitive - Suspected on Forests (LOLO) Species of Conservation Concern on Forests (FLAT)		3	Wetland/Riparian
			Species Occurrences verified in these Counties: Flathead, Lincoln, Powell State Rank Reason: Rare in Montana, where it is known from fens and wet meadows in the northwest corner of the state. Generally does not appear to be threatened by any particular activities, though populations are susceptible to hydrologic changes.						
Carex comosa Bristly Sedge		Cyperaceae Sedges	G5	S1S2				1	Wetland/Riparian
			Species Occurrences verified in these Counties: Flathead State Rank Reason: Only one known location in Montana on the shore of Flathead Lake. Occurrence is threatened by erosion caused by wave action and artificially high lake levels.						
Carex crawei Crawe's Sedge		Cyperaceae Sedges	G5	S2S3				3	Wetland/Riparian
			Species Occurrences verified in these Counties: Cascade, Pondera, Powell, Prairie, Teton State Rank Reason: Rare in Montana, where it is known from several areas. A few sites contain moderate to large populations. Trend data are lacking for the species. Negative impacts to populations from hydrologic changes are a potential threat.						
Carex glacialis Alpine Sedge		Cyperaceae Sedges	G5	S3				3	
			Species Occurrences verified in these Counties: Flathead, Lewis and Clark, Pondera State Rank Reason: <i>Carex glacialis</i> occurs throughout Canada, and has recently been discovered in the United States where it occurs at 4 locations in Montana. It grows in limestone fellfield habitats within the alpine. Populations are few, but appear stable. Surveys are needed to explore potential habitat, map its distribution, and determine population sizes.						
Carex gravida Heavy Sedge		Cyperaceae Sedges	G5	S3		Sensitive - Known on Forests (CG)		2	Wetland/Riparian
			Species Occurrences verified in these Counties: Big Horn, Carter, Fallon, McCone, Powder River, Richland, Rosebud State Rank Reason: <i>Carex gravida</i> has been found at a few widely scattered locations in eastern Montana, and is not generally abundant where it occurs. However, it is likely that the species is more abundant than the current data shows. Habitats include moist, green ash woodlands, which are attractive to livestock, and it may be particularly vulnerable to moderate grazing because of its caespitose growth form. These habitats are also quite vulnerable to invasion by non-native plants.						
Carex idahoensis Idaho Sedge	Carex parryana ssp. idahoensis	Cyperaceae Sedges	G3	S3		Sensitive - Known on Forests (BD)	SENSITIVE	2	Wetland/Riparian
			Species Occurrences verified in these Counties: Beaverhead, Broadwater, Deer Lodge, Gallatin, Madison, Powell, Silver Bow State Rank Reason: Idaho sedge is a regional endemic known from several dozen sites in Montana which cluster into approx 15-20 populations, most on public lands. The estimated number of stems is in the tens of thousands, but total occupied habitat has been estimated at less than 200 acres. The species is palatable, and populations may be affected by heavy grazing. Other risks are competition from exotic species, hydrologic alterations, agricultural development and road construction/maintenance. Updated population data and related site information are needed.						
Carex incurviformis Coastal Sand Sedge	Carex maritima var. incurviformis	Cyperaceae Sedges	G4G5	S2?				3	Wetland/Riparian
			Species Occurrences verified in these Counties: Deer Lodge, Glacier, Madison, Teton State Rank Reason: Five known occurrences in Montana, three are in Wilderness areas or Glacier National Park. However, all populations are apparently small to moderate in size based on limited survey data for the species. All occurrences are in alpine habitat that is not generally subject to human impacts.						
Carex lacustris Lake-bank Sedge		Cyperaceae Sedges	G5	S1S2		Species of Conservation Concern on Forests (FLAT)		3	Fens and marshes
			Species Occurrences verified in these Counties: Lake, Missoula State Rank Reason: A rare species in Montana, known only from a few occurrences from Lake County.						
Carex occidentalis Western Sedge		Cyperaceae Sedges	G4	S1				4	Dry, montane to alpine
			Species Occurrences verified in these Counties: Beaverhead, Silver Bow State Rank Reason: <i>Carex occidentalis</i> occurs on the northern edge of its range in Beaverhead County, Montana. This sedge is known from four locations of which specimens found at sites in 2018 and 1996 have been verified. The 1895 specimen is imprecisely mapped in Silver Bow County and may no longer be present and the 1984 report needs to be verified. Threats have not been identified and this species can be easily overlooked. Surveys that bring forth current data on locations, population sizes, habitat requirements, or threats is greatly needed.						
Carex petricosa Rock Sedge		Cyperaceae Sedges	G4	S1S2				3	Alpine
			Species Occurrences verified in these Counties: Glacier State Rank Reason: Rare in Montana, where it is currently known from one site in Glacier National Park. Very little data are available for the species in Montana. However, the potential for negative impacts to the populations appears to be low.						

Liparis loeselii Loesel's Twayblade		Orchidaceae Orchids	G5	S2		Species of Conservation Concern on Forests (FLAT)	3	Wetland/Riparian
Species Occurrences verified in these Counties: Lake State Rank Reason: Known from several occurrences clustered in a small area of the Swan Valley. Susceptible to changes in hydrology. May also be susceptible to impacts from fire.								
Muhlenbergia andina Foxtail Muhly		Poaceae Grasses	G4	S2S3			2	
Species Occurrences verified in these Counties: Broadwater, Carbon, Cascade, Gallatin, Granite, Lake, Lewis and Clark, Madison, Park, Sanders State Rank Reason: <i>Muhlenbergia andina</i> occurs widely scattered in western and south-central Montana. It grows in damp places, but often with well-drained soils. It can be found along streams, in wet meadows and seeps, and around hot springs. The low number of collections in combination with limited habitat and/or specific micro-habitat characteristics indicates it is either rare, declining, or over-looked in floristic surveys. Current data on locations, population sizes, habitat, and threats is greatly needed to better assess its status in Montana.								
Muhlenbergia minutissima Annual Muhly		Poaceae Grasses	G5	S3			4	
Species Occurrences verified in these Counties: Beaverhead, Gallatin, Madison, Missoula, Ravalli, Silver Bow State Rank Reason: <i>Muhlenbergia minutissima</i> is known from 7 locations observed from 1895 to 2015 in central and western Montana. It is also reported to occur in northeast Montana, but specimens have not been located (Peterson in FNA 2003). A 1941 occurrence near Belgrade has been searched for in recent decades, but not re-located (Matt Lavin personal communication). Plants can occupy disturbed areas, yet populations may not be persisting. Surveys that bring forth current data on locations, populations sizes, habitat requirements, or threats is needed.								
Najas guadalupensis Guadalupe Water-nymph		Najadaceae Water-nymph Family	G5	S2S3			4	Aquatic
Species Occurrences verified in these Counties: Carter, Cascade, Flathead, Lake, Lincoln, Ravalli, Sanders State Rank Reason: Rare. Currently documented from a few fresh water sites in the western and central portions of the state. Species is poorly documented in Montana and additional information on population levels, trends and threats is needed.								
Phippsia algida Ice Grass		Poaceae Grasses	G5	S2S3			3	Alpine
Species Occurrences verified in these Counties: Carbon, Stillwater State Rank Reason: Rare in Montana, where it has been documented from only a few sites on the Beartooth Plateau. Additional surveys of suitable habitat and revisits of documented occurrences are needed to more accurately assess the species' conservation status.								
Piperia elongata Dense-flower Rein Orchid	Habenaria elegans var. elata , Piperia elegans var. elata	Orchidaceae Orchids	G4	S1			4	
Species Occurrences verified in these Counties: Flathead, Missoula State Rank Reason: In Montana <i>Piperia elongata</i> is known from a single 1957 herbarium specimen collected in Lincoln County, and more recently from a few photographed specimens from Flathead, Lake, and Missoula Counties. However, the more recent observations lack data on population size and extent, habitat condition, threats, and other information. Surveys are needed to better document its status in Montana.								
Poa laxa ssp. banffiana Banff Bluegrass		Poaceae Grasses	G5?T2	S1			3	Alpine
Species Occurrences verified in these Counties: Glacier								
Potamogeton obtusifolius Blunt-leaved Pondweed		Potamogetonaceae Pondweeds	G5	S3		Sensitive - Suspected on Forests (LOLO) Species of Conservation Concern on Forests (HLC)	3	Aquatic
Species Occurrences verified in these Counties: Flathead, Glacier, Lake, Missoula, Powell State Rank Reason: Known from over a dozen occurrences in northwest Montana. Several contain moderate to large-size populations and occur in valley and foothill locations in a variety of federal, state, and private ownerships. A few populations are on lands managed specifically for their conservation value. Some populations are vulnerable to impacts associated with development, recreation and increased sediment and nutrient loads.								
Puccinellia lemmonii Lemmon's Alkaligrass		Poaceae Grasses	G4	S1S2			3	Wetland/Riparian
Species Occurrences verified in these Counties: Beaverhead State Rank Reason: Very rare in Montana where it is known only from Beaverhead County on BLM and State Trust Lands. At least one site is actively grazed, though its susceptibility and response to such activity is uncertain.								
Scheuchzeria palustris Pod Grass		Scheuchzeriaceae Pod-grasses	G5	S3		Sensitive - Known on Forests (BD, KOOT, LOLO) Sensitive - Suspected on Forests (BRT)	3	Wetland/Riparian

			Species Occurrences verified in these Counties: Flathead, Granite, Lake, Lincoln, Missoula State Rank Reason: Known in Montana from several dozen fens west of the Continental Divide. Several locations are known only from historical surveys or collections, or from sites that need additional surveys to document the populations. The majority of populations are on National Forest lands with MT State Trust lands, private and National Park lands supporting the remaining occurrences. Populations are primarily vulnerable to activities that change the hydrology of the occupied fen and wetland habitats.					
Schoenoplectus heterochaetus Slender Bulrush	Scirpus heterochaetus	Cyperaceae Sedges	G5	S1S2			4	Wetland/Riparian
			Species Occurrences verified in these Counties: Lake, Phillips State Rank Reason: Information on the species is lacking within Montana where it is recorded from only two poorly documented sites. However, its apparent rarity in the state warrants a high conservation status rank.					
Schoenoplectus subterminalis Water Bulrush	Scirpus subterminalis	Cyperaceae Sedges	G5	S3		Sensitive - Known on Forests (KOOT, LOLO) Species of Conservation Concern on Forests (HLC)	3	Wetland/Riparian
			Species Occurrences verified in these Counties: Flathead, Lake, Lewis and Clark, Lincoln, Missoula State Rank Reason: Over a dozen known occurrences in western Montana, most of which are moderate to large-sized populations primarily on National Forest lands. Populations are potentially vulnerable to changes in water levels or increases in nutrient and sediment loads associated with development, agriculture or adjacent timber harvesting.					
Scolochloa festucacea Sprangletop		Poaceae Grasses	G5	S1			4	
			Species Occurrences verified in these Counties: Flathead State Rank Reason: <i>Scolochloa festucacea</i> occurs through most of Canada and in portions of mid-western and western States. In Montana it is known from 3 locations collected from 1949 to 1999 in Flathead County. A fourth location from a specimen with a poorly defined location in Carbon county needs to be verified. Surveys to find this species have been unsuccessful.					
Sisyrinchium septentrionale Northern Blue-eyed-grass		Iridaceae Irises	G4	S1S2			4	Wetland/Riparian
			Species Occurrences verified in these Counties: Sheridan State Rank Reason: Rare in Montana, where it is known from one prairie site in the northeastern corner of the state. Population information and related habitat data from the known location are lacking.					
Spiranthes diluvialis Ute Ladies'-tresses	Ute Lady's-tresses, Ute Ladies-tresses	Orchidaceae Orchids	G2G3	S1S2	LT		1	Wetland/Riparian
			Species Occurrences verified in these Counties: Beaverhead, Broadwater, Gallatin, Jefferson, Madison State Rank Reason: <i>Spiranthes diluvialis</i> is known from a small number of occurrences in southwest and south-central Montana. Plants occur in the valleys of the Missouri, Jefferson, Beaverhead, Ruby, and Madison River drainages where it is restricted in area by specific hydrologic requirements. Many populations have less than 100 individuals, though a couple have over 500 plants. Sites are susceptible to hydrologic changes and weed invasion. Large areas of habitat have been converted to agricultural uses. Agricultural practices can hinder or promote plants depending upon their timing with critical reproductive stages. A few populations occur along highway right-of-ways. Most populations occur on private lands and only one occurrence is currently provided some potential protection or management for its conservation value.					
Sporobolus compositus Tall Dropseed	Sporobolus asper	Poaceae Grasses	G5	SH			4	Forests/Grasslands (open, plains)
			Species Occurrences verified in these Counties: Big Horn, Carter, Custer State Rank Reason: Known in Montana from 3 collections; a 1939 collection near Ekalaka, a 1957 collection from Fort Keogh Livestock and Range Laboratory and a 1980 collection from Bighorn County.					
Sporobolus neglectus Small Dropseed		Poaceae Grasses	G5	S1S2			4	Grasslands (low-elevation)
			Species Occurrences verified in these Counties: Gallatin, Sanders, Valley, Wheatland State Rank Reason: Rare in Montana, where it is known from a few widely scattered and poorly documented sites.					
Stipa lettermanii Letterman's Needlegrass	Achnatherum lettermanii	Poaceae Grasses	G5	S1S3		Species of Conservation Concern on Forests (HLC)	3	Talus and Grasslands (low-elevation)
			Species Occurrences verified in these Counties: Beaverhead, Big Horn, Carbon, Gallatin, Madison, Mineral, Park, Powell State Rank Reason: Documented from several locations in the southern portion of the state. However, population levels, site characteristics and related information needed to determine the species' status are lacking.					
Tofieldia pusilla Small Tofieldia		Liliaceae Lilies	G5	S2			3	Alpine
			Species Occurrences verified in these Counties: Flathead, Glacier State Rank Reason: Very rare in Montana, where it is known from only a very small area in Glacier National Park.					
Trichophorum alpinum Hudson's Bay Bulrush	Scirpus hudsonianus , Eriophorum alpinum	Cyperaceae Sedges	G5	S2		Species of Conservation Concern on Forests (FLAT)	3	Fens and cold, wet slopes

Fissidens fontanus Flat Pocket Moss	A Pocket Moss	Fissidentaceae	G5	S1				
			Species Occurrences verified in these Counties: Granite					
Grimmia brittoniae Britton's Dry Rock Moss	Britton's Black Rock Moss	Grimmiaceae	G2	S2		Sensitive - Known on Forests (KOOT, LOLO) Species of Conservation Concern on Forests (FLAT)		
			Species Occurrences verified in these Counties: Flathead, Sanders					
Grimmia incurva Curved Dry Rock Moss	Curved Black Rock Moss	Grimmiaceae	GNR	S1				
			Species Occurrences verified in these Counties: Ravalli					
Hamatocaulis vernicosus Hamatocaulis Moss	Drepanocladus vernicosus	Amblystegiaceae	G5	S1				
			Species Occurrences verified in these Counties: Flathead, Lake, Lincoln					
Haplodontium macrocarpum Waterfall Copper Moss	Mielichhoferia macrocarpa , Bryum porsildii	Bryaceae	G2G3	S1				
			Species Occurrences verified in these Counties: State Rank Reason: One specimen collected from a population growing on a wet limestone cliff in Park County, MT in 1973.					
Hennediella heimii Heim's Hennediella Moss	Desmatodon heimii	Pottiaceae	G5	S1				
			Species Occurrences verified in these Counties: Ravalli					
Homalothecium megaptilum Giant Golden Moss	Trachybryum megaptilum	Brachytheciaceae	G4	S1				
			Species Occurrences verified in these Counties: Lake, Lincoln, Mineral, Sanders State Rank Reason: Endemic to western North America. In Montana it occurs on the eastern edge of its distribution.					
Hygroamblystegium varium ssp. noterophilum A Conecap Moss	Hygroamblystegium noterophilum A Hygroamblystegium Moss	Amblystegiaceae	G5T4	S1				
			Species Occurrences verified in these Counties: Cascade, Granite					
Leucolepis acanthoneuron Umbrella Moss	Leucolepis menziesii	Mniaceae	G4G5	S1				
			Species Occurrences verified in these Counties: Lincoln, Sanders					
Meesia longiseta Meesia Moss		Meesiaceae	G5	S1				
			Species Occurrences verified in these Counties: Flathead, Park					
Meesia triquetra Meesia Moss		Meesiaceae	G5	S2		Sensitive - Known on Forests (BRT, CG, KOOT) Sensitive - Suspected on Forests (LOLO) Species of Conservation Concern on Forests (FLAT)		
			Species Occurrences verified in these Counties: Carbon, Flathead, Glacier, Lake, Lincoln, Ravalli, Sanders, Teton					
Meesia uliginosa Meesia Moss	Broad-leaved Hump Moss	Meesiaceae	G5	S1S2				
			Species Occurrences verified in these Counties: Flathead, Glacier, Lake, Lincoln, Sanders					
Meiotrichum lyallii Lyall's Polytrichum Moss	Polytrichum lyallii , Polytrichadelphus lyallii , Polytrichastrum lyallii	Polytrichaceae	G3G5	S1				
			Species Occurrences verified in these Counties: Flathead, Sanders					
Myurella tenerrima A Mousetail Moss		Pterigynandraceae	G5	S1				
			Species Occurrences verified in these Counties: Glacier					
Neckera douglasii Douglas' Neckera Moss		Neckeraceae	G4	S1				
			Species Occurrences verified in these Counties: Flathead, Lake, Sanders					
Paludella squarrosa Angled Paludella Moss		Meesiaceae	G5	S1S2				

			Species Occurrences verified in these Counties: Beaverhead, Carbon, Flathead, Glacier					
Paraleucobryum enerve A Windblown Moss		Dicranaceae	G5?	S1				
			Species Occurrences verified in these Counties: Flathead, Glacier, Stillwater					
Physcomitrium hookeri Hooker's Physcomitrium Moss		Funariaceae	G2G4	S1				
			Species Occurrences verified in these Counties: Ravalli, Roosevelt					
Porotrichum bigelovii Bigelow's Porotrichum Moss		Thamnobryaceae	G4	S1				
			Species Occurrences verified in these Counties: Ravalli					
Pseudocrossidium obtusulum A Pseudocrossidium Moss		Pottiaceae	GU	S1				
			Species Occurrences verified in these Counties: Musselshell, Ravalli					
Ptychostomum schleicheri Schleicher's Ptychostomum Moss	Bryum schleicheri	Bryaceae	G5?	S1				
			Species Occurrences verified in these Counties: Glacier					
Rhynchostegium aquaticum Aquatic Rhynchostegium Moss	Eurhynchium riparioides, Platyhypnidium riparioides, Platyhypnidium aquaticum	Brachytheciaceae	GNR	S1				
			Species Occurrences verified in these Counties: Lake, Lincoln, Sanders					
Sarmentypnum exannulatum Warnstorfia Moss	Warnstorfia exannulata	Amblystegiaceae	G5	S1				
			Species Occurrences verified in these Counties: Beaverhead, Carbon, Deer Lodge, Flathead, Glacier, Lincoln, Park, Sweet Grass					
Scorpidium revolvens Limprichtia Moss	Drepanocladus revolvens, Limprichtia revolvens	Amblystegiaceae	G5	S1				
			Species Occurrences verified in these Counties: Flathead, Gallatin, Glacier, Lake, Missoula, Sanders, Stillwater, Sweet Grass, Teton					
Scorpidium scorpioides A Scorpidium Moss		Amblystegiaceae	G5	S2		Sensitive - Known on Forests (KOOT) Species of Conservation Concern on Forests (FLAT, HLC)		
			Species Occurrences verified in these Counties: Flathead, Glacier, Lake, Lewis and Clark, Lincoln, Missoula, Powell, Teton					
Sphagnum angustifolium Narrowleaf Peatmoss		Sphagnaceae Peat Mosses	G5	S2				
			Species Occurrences verified in these Counties: Beaverhead, Flathead, Lincoln, Missoula, Park, Sanders					
Sphagnum centrale A Peatmoss		Sphagnaceae Peat Mosses	G5	S1				
			Species Occurrences verified in these Counties: Flathead, Missoula, Ravalli, Sanders					
Sphagnum compactum Cushion Peatmoss	Low Peatmoss	Sphagnaceae Peat Mosses	G5	S1				
			Species Occurrences verified in these Counties: Granite, Meagher					
Sphagnum contortum Contorted Sphagnum Moss		Sphagnaceae Peat Mosses	G5	S1				
			Species Occurrences verified in these Counties: Flathead, Lincoln					
Sphagnum fimbriatum Fringed Bogmoss	Ragged Hair Peatmoss	Sphagnaceae Peat Mosses	G5	S1		Species of Conservation Concern on Forests (HLC)		
			Species Occurrences verified in these Counties: Beaverhead, Flathead, Granite, Lewis and Clark, Park					
Sphagnum fuscum Brown Hair Peatmoss	Brown Peatmoss	Sphagnaceae Peat Mosses	G5	S2				
			Species Occurrences verified in these Counties: Carbon, Flathead, Lake, Lincoln, Missoula, Ravalli					
Sphagnum girgensohnii Star Hair Peatmoss	Girgensohn's Peatmoss	Sphagnaceae Peat Mosses	G5	S1				
			Species Occurrences verified in these Counties: Lincoln					
Sphagnum magellanicum Red Spoon Peatmoss	Magellan's Peatmoss	Sphagnaceae Peat Mosses	G5	S1		Species of Conservation Concern on Forests (FLAT)		

Dactylina ramulosa Frosted Finger Lichen		Parmeliaceae	G5	S2				
			Species Occurrences verified in these Counties: Park, Ravalli					
Gyalectaria diluta Diluted Wart Lichen	Pertusaria diluta	Coccotremataceae	GNR	S1				
			Species Occurrences verified in these Counties: State Rank Reason: This species was first recognized in Montana. The Type specimen is from the Cabinet Mountains and is currently the only Montana occurrence.					
Lobaria amplissima Large Lungwort Lichen		Lobariaceae	GNR	SNR				
			Species Occurrences verified in these Counties: State Rank Reason: Known from one location in western Montana.					
Lobaria anomala Netted Lungwort Lichen	Pseudocyphellaria anomala	Lobariaceae	G2G4	S1				
			Species Occurrences verified in these Counties: Lake State Rank Reason: Known in western Montana from a few locations.					
Lobaria hallii Gray Lungwort Lichen		Lobariaceae	G4?	S2				
			Species Occurrences verified in these Counties: Flathead, Lake, Lincoln, Missoula, Sanders State Rank Reason: Known from several locations in western Montana.					
Lobaria linita Cabbage Lungwort Lichen		Lobariaceae	G5	S1				
			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Known from very few locations in western Montana.					
Lobaria scrobiculata Textured Lungwort Lichen		Lobariaceae	G5	S1				
			Species Occurrences verified in these Counties: Lake, Mineral State Rank Reason: Known from one location in western Montana.					
Melanohalea septentrionalis Northern Camouflage Lichen		Parmeliaceae	G5	S1				
			Species Occurrences verified in these Counties: State Rank Reason: Montana occurs on the southern edge of this species range, where it has been found occasionally.					
Nodobryoria subdivergens Alpine Foxtail Lichen	Alectoria subdivergens, Bryoria subdivergens	Parmeliaceae	G2G3	S1S2		Sensitive - Known on Forests (BRT, KOOT)		
			Species Occurrences verified in these Counties: Glacier, Lincoln, Ravalli State Rank Reason: Known from several locations in western Montana where its abundance is always sparse.					
Normandina pulchella Elf-Ear Lichen		Verrucariaceae	G4G5	S1				
			Species Occurrences verified in these Counties: Missoula, Ravalli State Rank Reason: In the Rocky Mountains, this lichen has a spotty distribution. Known in Montana from one location.					
Parmeliella triptophylla Fingered Shingle Lichen	Pannaria triptophylla Black-bordered Shingle Lichen	Pannariaceae	G5	S1				
			Species Occurrences verified in these Counties: Glacier, Lake, Missoula, Ravalli State Rank Reason: Locally rare when found.					
Peltigera gowardii Western Waterfan Lichen	Peltigera hydrothyria [name misapplied in western North America], Hydrothyria venosa	Peltigeraceae	G3G4	S1				
			Species Occurrences verified in these Counties: Missoula, Ravalli State Rank Reason: Known from a few sites in western Montana.					
Peltigera pacifica Fringed Pelt Lichen		Peltigeraceae	G3G4	S1				
			Species Occurrences verified in these Counties: State Rank Reason: Known from one location in western Montana, but expected to be more present.					
Phaeophyscia kairamoi Least Shadow Lichen		Physciaceae	G4G5	S2				
			Species Occurrences verified in these Counties: Flathead, Lake State Rank Reason: This species occurs sporadically in the northern United States and southern Canada and is known from a few locations in western Montana.					
Ramalina labiosorediata Chalky Bush Lichen	Ramalina pollinaria	Ramalinaceae	G4	S1				
			Species Occurrences verified in these Counties: Lake State Rank Reason: Known in western Montana from several locations.					
Ramalina obtusata Hooded Bush Lichen		Ramalinaceae	G5	S2				
			Species Occurrences verified in these Counties: Flathead, Lake, Ravalli State Rank Reason: In Montana sporadic occurrences have been found in western Montana.					

Rhizoplaca haydenii Hayden's Rimmed Navel Lichen		Lecanoraceae	G2G3	S1S2				
			Species Occurrences verified in these Counties: Beaverhead, Carbon State Rank Reason: Known from a few locations in south-central to southeastern Montana. This species is also likely to be found in appropriate habitats in southwestern Montana. Both subspecies are found in Montana: <i>R. haydenii</i> ssp. <i>haydenii</i> and <i>R. haydenii</i> ssp. <i>arbuscular</i> .					
Sclerophora amabilis Lovely Pin Lichen		Coniocybaceae	G4G5	S1				
			Species Occurrences verified in these Counties: Lincoln State Rank Reason: In Montana known from one location.					
Solorina bispora Lesser Tundra Owl Lichen		Peltigeraceae	G5	S1S2				
			Species Occurrences verified in these Counties: Beaverhead, Carbon, Flathead, Glacier, Missoula State Rank Reason: Known from a few locations in western Montana.					
Solorina octospora Greater Tundra Owl Lichen		Peltigeraceae	G3G5	S1				
			Species Occurrences verified in these Counties: State Rank Reason: In Montana known from one location in the northwest.					
Solorina spongiosa Fringed Chocolate Chip Lichen		Peltigeraceae	G4G5	S1S2				
			Species Occurrences verified in these Counties: Flathead, Lake, Lewis and Clark State Rank Reason: Known from a few locations in western and central portions of Montana.					
Sphaerophorus tuckermanii Tuckermann's Coral Lichen		Sphaerophoraceae	G5	S1				
			Species Occurrences verified in these Counties: State Rank Reason: Known from two locations in northwestern Montana.					
Stereocaulon paschale Easter Foam Lichen		Stereocaulaceae	G5	S1S2				
			Species Occurrences verified in these Counties: Lake State Rank Reason: Known from a few locations in northwest and south-central Montana.					
Umbilicaria hirsuta Granulating Rocktripe Lichen		Umbilicariaceae	G2G4	S1				
			Species Occurrences verified in these Counties: State Rank Reason: This species is apparently rare throughout its range in North America. In Montana it is known from one location.					
Verrucaria kootenaica Kootenai Speck Lichen		Verrucariaceae	G2	S1S2				
			Species Occurrences verified in these Counties: Flathead, Lake State Rank Reason: Known in western Montana from a few locations.					

All Records (no filtering)

FLOWERING PLANTS - DICOTS (MAGNOLIOPSIDA)								53 SPECIES	
SCIENTIFIC NAME COMMON NAME TAXA SORT	OTHER NAMES	FAMILY (SCIENTIFIC) FAMILY (COMMON)	GLOBAL RANK	STATE RANK	USFWS	USFS	BLM	MNPS THREAT CATEGORY	HABITAT
Agoseris lackschewitzii Pink Agoseris	Agoseris aurantiaca var. aurantiaca , Agoseris carnea	Asteraceae Aster/Sunflowers	G4Q	S3S4					
			Species Occurrences verified in these Counties: Beaverhead, Carbon, Cascade, Deer Lodge, Gallatin, Granite, Judith Basin, Madison, Meagher, Park, Silver Bow, Sweet Grass State Rank Reason: See rank details.						
Allotropa virgata Candystick		Ericaceae Heath Family	G4	S3S4					
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Granite, Ravalli State Rank Reason: Limited distribution and small population sizes make the species potentially vulnerable to impacts to its habitat, primary lodgepole pine stands. Trend and monitoring data for the species are lacking. However, populations are presumed to be relatively stable at the present time.						
Aquilegia jonesii Jones' Columbine		Ranunculaceae Buttercup Family	G3	S3S4					
			Species Occurrences verified in these Counties: Cascade, Fergus, Flathead, Glacier, Judith Basin, Lewis and Clark, Meagher, Park, Stillwater, Sweet Grass, Teton						
Arabidopsis lyrata Lyre-leaf Rockcress	Arabis lyrata , Arabis kamchatica	Brassicaceae Mustards	G5	SH					NA
			Species Occurrences verified in these Counties: Flathead State Rank Reason: Known from one 1952 collection near Mount Brown in Glacier National Park.						
Atriplex canescens Four-wing Saltbush		Amaranthaceae Amaranth (Pigweed) Family	G5	S3S4					
			Species Occurrences verified in these Counties: Carter, Park, Richland, Silver Bow, Valley						
Atriplex suckleyi Suckley's Saltbush	Atriplex dioica (Nutt.) Macbr. [not Raf.], Endolepis dioica	Amaranthaceae Amaranth (Pigweed) Family	G4	S3S4					
			Species Occurrences verified in these Counties: Carbon, Carter, Fallon, Phillips, Powder River, Richland, Toole, Valley, Wheatland State Rank Reason: Few collections from Montana, mostly along the Missouri River Breaks. However, this species has weedy tendencies.						
			MONT collections from Valley, McCone Counties.						

Eutrema salsugineum Saltwater Cress	Arabidopsis salsuginea, Thellungiella salsuginea	Brassicaceae Mustards	G5?	SH				
			Species Occurrences verified in these Counties:					
Gaultheria ovatifolia Slender Wintergreen		Ericaceae Heath Family	G5	S3S4				
			Species Occurrences verified in these Counties: Flathead, Glacier, Lake, Lincoln, Mineral, Sanders					
Geocaulon lividum Northern Toadflax	Comandra lividum	Santalaceae Sandalwood Family	G5	S3S4				
			Species Occurrences verified in these Counties: Flathead, Lake, Lincoln, Missoula, Sanders					
Gilia tweedyi Tweedy's Gilia	Gilia sinuata var. tweedyi, Gilia inconspicua var. tweedyi	Polemoniaceae Phlox Family	G4G5Q	S3S4				
			Species Occurrences verified in these Counties: Beaverhead, Carbon State Rank Reason: Gilia tweedyi is locally common on the south and west sides of the Pryor Mountains in the drainages of the Bighorn and Clarks Fork of the Yellowstone rivers and is also known from Beaverhead County.					
Hedysarum alpinum Alpine Sweet-vetch		Fabaceae Pea Family	G5	S3S4				
			Species Occurrences verified in these Counties: Flathead, Gallatin, Glacier, Lake, Missoula, Phillips					
Hymenoxys torreyana Torrey Bitterweed	Tetaneuris torreyana	Asteraceae Aster/Sunflowers	G4	S3S4				
			Species Occurrences verified in these Counties: Carbon					
Impatiens ecalcarata Spurless Touch-me-not		Balsaminaceae Impatiens	G3G4	S3S4				
			Species Occurrences verified in these Counties: Gallatin, Lake, Missoula, Sanders					
Linanthastrum nuttallii Nuttall's Linanthus	Linanthus nuttallii, Leptosiphon nuttallii	Polemoniaceae Phlox Family	G5	S3S4				
			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Reported as locally common in the Bitterroot Mountains by Lesica & Shelly (1991).					
Lomatium bicolor Bicolor Biscuitroot		Apiaceae Parsley/Carrot Family	G4	S3S4				
			Species Occurrences verified in these Counties: Glacier, Ravalli					
Lorandersonia linifolia Spearleaf Rabbitbrush	Chrysothamnus viscidiflorus var. linifolius, Chrysothamnus linifolius	Asteraceae Aster/Sunflowers	G5	S3S4				
			Species Occurrences verified in these Counties: Custer, Powder River					
Madia minima Small-headed Tarweed	Hemizonella minima	Asteraceae Aster/Sunflowers	G4	S3S4				
			Species Occurrences verified in these Counties: Flathead, Granite, Lincoln, Missoula, Ravalli, Sanders					
Mimulus suksdorfii Suksdorf Monkeyflower		Phrymaceae Lopseed Family	G4	S3S4				
			Species Occurrences verified in these Counties: Beaverhead, Carbon, Gallatin, Lewis and Clark, Madison, Missoula, Park, Rosebud, Silver Bow					
Musineon vaginatum Rydberg's Parsley		Apiaceae Parsley/Carrot Family	G3G4	S3S4				
			Species Occurrences verified in these Counties: Big Horn, Carbon, Gallatin, Granite, Lewis and Clark, Missoula, Rosebud, Teton State Rank Reason: See rank details.					
Orobanche corymbosa Flat-topped Broomrape		Orobanchaceae Broomrape Family	G4	S3S4				
			Species Occurrences verified in these Counties: Beaverhead, Deer Lodge, Granite, Jefferson, Madison, Powell, Ravalli					
Oxytropis lagopus var. conjugans Hare's-foot Locoweed		Fabaceae Pea Family	G4G5T3T4	S3S4			3	Sagebrush (low-elevation)
			Species Occurrences verified in these Counties: Beaverhead, Broadwater, Granite, Jefferson, Lewis and Clark, Meagher, Pondera, Powell, Teton State Rank Reason: See rank details.					
Pedicularis oederi Oeder's Lousewort		Orobanchaceae Broomrape Family	G5	S3S4				
			Species Occurrences verified in these Counties: Carbon, Stillwater					
Pediomelum hypogaeum var. hypogaeum Little Indian Breadroot	Psoralea hypogaea	Fabaceae Pea Family	G5T4	S3S4			3	Grasslands/Woodlands (Open, sandy soil)
			Species Occurrences verified in these Counties: Carter, Cascade, Chouteau, Fergus, Golden Valley, Petroleum, Powder River, Rosebud State Rank Reason: <i>Pediomelum hypogaeum</i> was first documented in Montana from a specimen collected in 1886 in Cascade County (F.W. Anderson (s.n.) NY). Surveys in the 1980s and 1990s resulted in collections made from Fergus, Petroleum, and Rosebud Counties (http://rmh.uwyo.edu and https://www.pnwherbaria.org/). Additional observations found in other counties of southeast Montana should be verified. <i>Pediomelum hypogaeum</i> is categorized as a Potential Species of Concern because populations tend to be smaller and widely spaced, habitat may be limited, and its biology may make it more vulnerable to extirpation or just slow to recover from adverse impacts. Current information on plant locations, population sizes, and threats is needed. See rank details.					
Penstemon laricifolius Larch-leaf Beardtongue		Plantaginaceae Plantain Family	G4	S3S4				
			Species Occurrences verified in these Counties: Big Horn, Carbon State Rank Reason: In Montana, <i>Penstemon laricifolius</i> is known from Carbon County where it is common on the south and west flanks of the Pryor Mountains.					
Phacelia scopulina Dwarf Phacelia	Phacelia lutea var. scopulina	Hydrophyllaceae Waterleaf Family	G4	SH				Alkaline sites

			Species Occurrences verified in these Counties: Beaverhead, Madison, Silver Bow State Rank Reason: Known in Montana from one 1885 collection by P.A. Rydberg near Melrose, probably in Silver Bow County.					
Phlox andicola Plains Phlox		Polemoniaceae Phlox Family	G4	S3S4			3	Open sites (Sand to clay soils)
			Species Occurrences verified in these Counties: Carter, Dawson, Phillips, Powder River, Rosebud, Sheridan State Rank Reason: Plains phlox reaches the western margin of its range in Montana's eastern counties. It has been documented from relatively few locations, but surveys during its early blooming season have been few, and additional spring inventory work may locate more populations. It likely tolerates grazing and may benefit from some level of disturbance.					
Polygonum austini Austin's Knotweed	Polygonum douglasii ssp. austini	Polygonaceae Buckwheat Family	G5T4	S3S4		Sensitive - Known on Forests (BD) Sensitive - Suspected on Forests (CG) Species of Conservation Concern on Forests (HLC)	2	Rock/Talus
			Species Occurrences verified in these Counties: Broadwater, Flathead, Glacier, Granite, Lewis and Clark, Madison, Meagher, Park, Pondera, Powell, Teton State Rank Reason: Austin's knotweed is sparsely distributed in mountainous areas of Montana from the Rocky Mountain Front to the Madison and Gallatin Ranges. Sites are usually on open, gravelly, sparsely-vegetated slopes with shale-derived soils and as such are not generally impacted by human activity. Some sites however, are along forest roads and are susceptible to weed invasion and other disturbances. The probability of finding additional occurrences appears to be good since large areas of suitable habitat across western and central Montana remain unsurveyed for the species.					
Ranunculus hyperboreus High Northern Buttercup	Ranunculus natans	Ranunculaceae Buttercup Family	G5	S3S4				Wetland/Riparian (Montane)
			Species Occurrences verified in these Counties: Beaverhead, Carbon, Deer Lodge, Gallatin, Jefferson, Madison, Missoula, Silver Bow, Valley State Rank Reason: Known from several southwest and south-central counties in Montana. See rank details for additional information.					
Sedum borschii Borsch's Stonecrop	Sedum leibergii	Crassulaceae Stonecrops	G3?	S3S4				
			Species Occurrences verified in these Counties: Beaverhead					
Solidago velutina Three-nerved Goldenrod	Solidago sparsiflora	Asteraceae Aster/Sunflowers	G5?	SH				NA
			Species Occurrences verified in these Counties: State Rank Reason: Few-flowered goldenrod is known in Montana from 1 specimen collection from the Stillwater River Valley, which lacks precise locality data. Other reports of this species from the state are based on mis-identified specimens. Additional data are needed.					
Sphaeralcea munroana White-stemmed globemallow		Malvaceae Mallow Family	G4	S3S4			3	Sagebrush-Grasslands (low-elevation)
			Species Occurrences verified in these Counties: Beaverhead, Jefferson, Park State Rank Reason: Peripheral in southwest Montana where it is known from a few locations. Additional survey and monitoring data are needed. Most documented locations are along roads and 2-tracks, as such, at least several of the populations may be adventive or introduced. Species appears to be tolerant of or perhaps benefits from some disturbance activity. Additional information concerning the conservation needs and population dynamics of this species in Montana is needed to clarify its status.					
Stanleya tomentosa Woolly Prince's plume		Brassicaceae Mustards	G4	S3S4				
			Species Occurrences verified in these Counties: Beaverhead, Carbon State Rank Reason: See rank details.					
Stanleya viridiflora Green Prince's plume		Brassicaceae Mustards	G4	S3S4				
			Species Occurrences verified in these Counties: Beaverhead, Madison State Rank Reason: See rank details.					
Stenotus multicaulis Many-stem Goldenweed	Oonopsis multicaulis, Haplopappus multicaulis	Asteraceae Aster/Sunflowers	G4	S3S4				
			Species Occurrences verified in these Counties: Carbon, Carter, Fallon State Rank Reason: Though restricted in distribution in Montana to Carter County, it is common in some habitats, including along some roadsides at least on BLM lands. No apparent, substantial threats to the species' viability in the state exist.					
Streptanthella longirostris Streptanthella		Brassicaceae Mustards	G5	S3S4				
			Species Occurrences verified in these Counties: Carbon State Rank Reason: Uncommon in Montana and restricted in distribution to Carbon County. Population sizes are poorly documented and associated information on trends and threats are also lacking.					
Synthyris missurica Western Mountain kittentails		Plantaginaceae Plantain Family	G4	S3S4				
			Species Occurrences verified in these Counties: Ravalli State Rank Reason: Uncommon in Montana and restricted in distribution to the Bitterroot Mtns. Population sizes are poorly documented and associated information on trends and threats are also lacking.					
Tonestus pygmaeus Pygmy Goldenweed	Haplopappus pygmaeus	Asteraceae Aster/Sunflowers	G4	SH				Alpine
			Species Occurrences verified in these Counties: State Rank Reason: Known in Montana from 1 historical collection from Lolo Peak. Other historical locations previously reported for MT have all been based on mis-identified specimens of <i>Tonestus lyallii</i>					
Townsendia spatulata Sword Townsend-daisy		Asteraceae Aster/Sunflowers	G3	S3S4			3	

Callicladium haldanianum Pretty Branch Moss		Hypnaceae	G5	SH				
			Species Occurrences verified in these Counties: Flathead					
Calliergon richardsonii Richardson's Calliergon Moss		Amblystegiaceae	G5	SH				
			Species Occurrences verified in these Counties: Glacier, Park					
Dendroalsia abietina A Dendroalsia Moss		Leucodontaceae	G4	SH				
			Species Occurrences verified in these Counties:					
Dicranum fragilifolium Fragile Leaf Dicranum Moss		Dicranaceae	G4G5	SH				
			Species Occurrences verified in these Counties: Flathead, Glacier, Lake					
Dicranum spadiceum A Dicranum Moss	Dicranum angustum	Dicranaceae	G5	SNR				
			Species Occurrences verified in these Counties: Flathead, Glacier State Rank Reason: MT Botanist Mincemoyer downgraded species from S1 to SH for lack of knowledge of specimens after 1972 and was not aware of specimens collected in 1994 from Glacier NP and 1995 from Pine Butte.					
Distichium inclinatum Incline Thread Moss	Incline Distichium Moss	Ditrichaceae	G5	SH				
			Species Occurrences verified in these Counties: Glacier, Teton					
Entosthodon rubiginosus Rusty Cord Moss	Entosthodon Moss	Funariaceae	G1G3	SH				
			Species Occurrences verified in these Counties: Cascade					
Grimmia mollis A Dry Rock Moss	Hydrogrimmia mollis A Black Rock Moss	Grimmiaceae	G5	SH				
			Species Occurrences verified in these Counties: Flathead, Glacier					
Hygrohypnum cochlearifolium Ear-leaf Boat Moss	Ear-leaf Hygrohypnum Moss	Amblystegiaceae	G4	SH				
			Species Occurrences verified in these Counties: Lincoln					
Plagiobryum zieri Zierian Hump-Moss		Bryaceae	G5	SH				
			Species Occurrences verified in these Counties:					
Pseudocalliergon trifarium Blunt Water Moss	Calliergon trifarium Worm Moss	Amblystegiaceae	G5	SH				
			Species Occurrences verified in these Counties: Flathead, Glacier, Missoula					
Pseudocalliergon turgescens A Pseudocalliergon Moss	Scorpidium turgescens, Calliergon turgescens	Amblystegiaceae	G5	SH				
			Species Occurrences verified in these Counties: Flathead, Glacier, Teton					
Sarmentypnum sarmentosum A Sarmenthypnum Moss	Calliergon sarmentosum	Amblystegiaceae	G5	SNR				
			Species Occurrences verified in these Counties: Flathead, Glacier					
Tayloria acuminata Acuminate Dung Moss		Splachnaceae	G3G4	SH				
			Species Occurrences verified in these Counties:					
Thamnobryum neckeroides Necker's Thamnobryum Moss	A Tree Moss	Thamnobryaceae	G4	SH				
			Species Occurrences verified in these Counties: Flathead, Lake, Missoula, Sanders					
Tortula cernua A Tortella Moss	Desmatodon cernuus	Pottiaceae	G4G5	SH				
			Species Occurrences verified in these Counties:					

LICHENS (FUNGI)									8 SPECIES
SCIENTIFIC NAME COMMON NAME TAXA SORT	OTHER NAMES	FAMILY (SCIENTIFIC) FAMILY (COMMON)	GLOBAL RANK	STATE RANK	USFWS	USFS	BLM	MNPS THREAT CATEGORY	HABITAT
Brigantiaea praetermissa Brick-Spored Firedot Lichen		Brigantiaeaceae (Brigantiaeaceae)	GNR	S2S3					
			Species Occurrences verified in these Counties: Lake State Rank Reason: The type specimen is from Sanders County. This lichen is considered uncommon in western Montana and widely scattered in the Pacific Northwest.						
Cetraria sepincola Chestnut Wrinkled Lichen	Tuckermannopsis sepincola	Parmeliaceae	G5	S2S3					
			Species Occurrences verified in these Counties: Flathead, Lake, Madison, Mineral State Rank Reason: Known from many locations, associated with bogs, in western Montana.						

Evernia divaricata Mountain Oakmoss Lichen		Parmeliaceae	G4G5	S1S2				
			Species Occurrences verified in these Counties: Carbon, Lake, Missoula State Rank Reason: Populations have a very spotty distribution in Montana.					
Parmelia fraudans Pea-green Shield Lichen		Parmeliaceae	G5	S1				
			Species Occurrences verified in these Counties: State Rank Reason: Rare in the Pacific Northwest (McCune and Goward 2009); Infrequently collected in Montana and adjacent states.					
Platismatia herrei Tattered Rag Lichen		Parmeliaceae	G5	S1				
			Species Occurrences verified in these Counties: State Rank Reason: Known from a few locations in northwestern Montana.					
Platismatia stenophylla Ribbon Rag Lichen		Parmeliaceae	G5	S1				
			Species Occurrences verified in these Counties: Lake, Ravalli State Rank Reason: Known from a few locations in western Montana.					
Psora rubiformis Pea-green Scale Lichen		Psoraceae	G3G5	S1S2				
			Species Occurrences verified in these Counties: Flathead, Glacier, Lake, Madison, Rosebud State Rank Reason: In Montana widely scattered populations have been found in northwest, southwest, and southeast.					
Umbilicaria havaasii Havaas' Rocktripe Lichen		Umbilicariaceae	G4	S1				
			Species Occurrences verified in these Counties: Flathead, Ravalli State Rank Reason: Known from a few locations in western Montana. Montana occurs on the eastern edge of this species range.					

Special Status Species

0 Species

All Records (no filtering)

This section is not Filtered

ADDITIONS TO STATEWIDE LIST		
SPECIES	DATE	NOTES
Amorpha canescens Lead Plant	9/25/2021	<i>Amorpha canescens</i> was documented in 1922 from Powder River County (Lockhart 25, USFS-RM) and in 1948 from Carter County (Booth 2675, MONT). From 1983 to 2013 various field projects reported another 8 locations of <i>Amorpha canescens</i> , but provided no specimens or photographs to validate the identifications. A 1984 search to re-locate the plants found at the 1948 location was unsuccessful. In 2019 surveys on the Custer-Gallatin National Forest found and verified 10 sites in Montana (Hansen 196 and 264, MONTU; Hansen 2019). The 2019 observations found healthy, reproductive plants with no apparent threats. Relative to the State of Montana <i>Amorpha canescens</i> is ranked as a Species of Concern because it occupies relatively little habitat and almost half of the reported observations need to be validated before re-assessing its state status.
Elymus triticoides Beardless Wildrye	10/20/2020	<i>Elymus triticoides</i> occurs throughout the western United States but in Montana has only been documented in four western counties (Barkworth in Flora of North America 2007; revised draft treatment in the Manual of Montana Vascular Plants). This grass is known from fewer than five locations which are widely scattered. Plants can be confused with <i>Elymus smithii</i> and/or may be overlooked in our state. Surveys that accurately identify this grass and bring forth information on locations, population sizes, habitat conditions, and threats is greatly needed.
Calochortus bruneaunis Bruneau Mariposa Lily	9/23/2020	
Lemna valdiviana Pale Duckweed	9/8/2020	
Viola pedatifida Prairie Violet	9/1/2020	
Tetradymia spinosa Short-spine Horsebrush	9/1/2020	
Pinguicula macroceras California Butterwort	8/14/2020	
Eleocharis bella Delicate Spikerush	8/7/2020	
Amphiscirpus nevadensis Nevada Clubrush	7/10/2020	
Carex occidentalis Western Sedge	2/25/2020	
Myriophyllum quitense Andean Water-milfoil	10/31/2019	<i>Myriophyllum quitense</i> is an aquatic plant that recently (2008-2016) has been found in three waterbodies of Montana. Plants are found in slow-moving rivers that vary in water quality from the Madison River in Yellowstone National Park to Toston Reservoir on the Missouri River. These locations represent a very narrow geographical portion of Montana. Proper identification of <i>Myriophyllum</i> species require careful collections that obtain flowering or fruiting structures, use of an appropriate and current taxonomic key, and time spent studying the specimen. More surveys are greatly needed to assess the abundance and distribution of <i>Myriophyllum quitense</i> in Montana.
Navarretia divaricata Divaricate Navarretia	10/31/2019	<i>Navarretia divaricata</i> in Lesica et al. (2012) is based on a 1981 herbarium specimen (MONT 68910) collected in a pasture in Sanders County that was re-determined independently by Leigh Johnson (author for the <i>Navarretia</i> treatment for Flora of North America) and Matt Lavin (MONT curator) to be <i>Navarretia squarrosa</i> . In October 2019, retired USFS Botanist Craig Odegard brought to the MONTU herbarium his 2017 collection of <i>Navarretia divaricata</i> which came from a different location in Sanders County and has been verified by Shannon Kimball (MONTU Curator).
Muhlenbergia minutissima Annual Muhly	10/31/2019	<i>Muhlenbergia minutissima</i> is known from 7 locations observed from 1895 to 2015 in central and western Montana. It is also reported to occur in northeast Montana, but specimens have not been located (Peterson in FNA 2003). A 1941 occurrence near Belgrade has been searched for in recent decades, but not re-located (Matt Lavin personal communication). Plants can occupy disturbed areas, yet populations may not be persisting. Surveys that bring forth current data on locations, populations sizes, habitat requirements, or threats is needed.
Muhlenbergia andina Foxtail Muhly	10/31/2019	<i>Muhlenbergia andina</i> occurs widely scattered in western and south-central Montana. It grows in damp places, but often with well-drained soils. It can be found along streams, in wet meadows and seeps, and around hot springs. The low number of collections in combination with limited habitat and/or specific micro-habitat characteristics indicates it is either rare, declining, or over-looked in floristic surveys. Current data on locations, population sizes, habitat, and threats is greatly needed to better assess its status in Montana.
Dichanthelium acuminatum Panic Grass	10/31/2019	<i>Dichanthelium acuminatum</i> is common and ubiquitous in most of the U.S. and Canada (Freckmann and Lelong in FNA 2007). The species is polymorphic and 10 major subspecies have been described, but many overlap in characteristics and widespread introgression from other <i>Dichanthium</i> species contributes to taxonomic difficulties (Freckmann and Lelong in FNA 2007). However, only subspecies <i>sericeum</i> has been documented in Montana. <i>Dichanthelium acuminatum</i> ssp. <i>sericeum</i> colonizes wet soils around the edges of hot springs. It occurs widely scattered through south-central, southwest, and northwest Montana, where it can be locally common. Observation data is aging, and some re-visits to known populations did not re-locate the grass. Given its narrow habitat requirements, potential threats from ground disturbance and recreation, and lack of current data a Species of Concern rank is warranted. Current data on locations, population sizes, threats, and how it responds to natural and manmade disturbances are greatly needed.
Isoetes howellii Howell's Quillwort	9/25/2018	<i>Isoetes howellii</i> is known from about 5 locations in Northwestern Montana. Based on limited information threats appear to be minimal, but survey work to document locations, population sizes, and threats is greatly needed.
Isoetes echinospora Spiny-spore Quillwort	9/25/2018	<i>Isoetes echinospora</i> is known from 8 occurrences scattered in western Montana. At one occurrence, the species has been observed in 1940, 1967, and 1998 indicating persistence. However, current survey work is need to document locations, population sizes, and threats.
Isoetes occidentalis Western Quillwort	9/25/2018	<i>Isoetes occidentalis</i> is known from two locations in northwest Montana. Survey work to identify other locations, document population sizes, and determine threats is greatly needed.
Celastrus scandens Bittersweet	9/25/2018	<i>Celastrus scandens</i> occurs frequently in woodlands, rocky hillsides, thickets, fence rows, and roadsides in the Great Plains (McGregor 1986). The previous SH rank in Montana was based on a vague location provided on a 1975 herbarium specimen. In recent years it has been collected at four locations in woody draws. It appears that the Montana sites represent the western edge of its range, and currently it ranks as an S1. Additional surveys of woody draws are needed to accurately document its distribution and population size in Montana.

ADDITIONS TO STATEWIDE LIST		
SPECIES	DATE	NOTES
Astragalus ceramicus var. filifolius Painted Milkvetch	9/25/2018	Astragalus ceramicus variety filifolius is associated with sandy soils of the sandhills and sandstone outcrops in eastern Montana. It is known from about 20 occurrences observed mostly from 1983 to 2000. Some populations occur in State Parks, and current data on population sizes and threats is needed. The Flora of the Great Plains (1986) considered it rare in the Great Plains except for the Nebraska sandhill region where it was somewhat common. Based on aging data, limited distribution, and an association to specific habitat types it is considered a Species of Concern.
Impatiens aurella Pale-yellow Jewel-weed	9/25/2018	Impatiens aurella is known from about 20 locations documented from 1886 to 2016. It is consider uncommon in Lake and Flathead Counties, where the majority of observations have been found, and rare in other counties of western Montana. It grows in wet, often organic soil in both disturbed and undisturbed wetlands, and rarely appears abundant. However, it may require or persist better with some hydrological disturbance. Re-visits to known locations and more surveys are needed to better document locations, population sizes, and threats.
Astragalus ceramicus Pottery Milkvetch	9/25/2018	Astragalus ceramicus variety filifolius is associated with sandy soils of the sandhills and sandstone outcrops in eastern Montana. It is known from about 20 occurrences observed mostly from 1983 to 2000. Some populations occur in State Parks, and current data on population sizes and threats is needed. The Flora of the Great Plains (1986) considered it rare in the Great Plains except for the Nebraska sandhill region where it was somewhat common. Based on aging data, limited distribution, and an association to specific habitat types it is considered a Species of Concern.
Artemisia tilesii Tlesius Wormwood	9/25/2018	Artemisia tilesii is known from seven locations located at higher elevations in western Montana. The species can be difficult to separate from Artemisia ludoviciana and A. michauxiana. Survey work to identify occurrences, determine population sizes, and assess threats is greatly needed before re-evaluating its status.
Carex amplifolia Big-leaf Sedge	9/25/2018	Carex amplifolia occurs in temperate western North America where it is usually uncommon or rare from coastal lowlands to middle elevations in the mountains (FNA 2002). The previous SH rank in Montana was based on a 1978 herbarium specimen. In recent years it has been collected from several wetlands in Sanders and Flathead Counties. Additional wetland surveys are needed to accurately document its distribution and population size in Montana.
Cryptogramma cascadenis Cascade Rockbrake	9/27/2017	Cryptogramma cascadenis is known from 11 locations in western Montana, of which 2 locations are poorly defined and considered historical, 5 locations occur in Wilderness areas, and the remaining 4 locations occur on U.S. Forest Service lands. Although the fern is thought to be undercollected and could be more common, current population and location data is needed to remove this plant from the Species of Concern list.
Marsilea oligospora Pepperwort	9/27/2017	Marsilea oligospora has relatively recently been segregated from Marsilea vestita (FNA 1993). It is quite common around Ninepipes National Wildlife Refuge, but has not been documented elsewhere in Montana. Observation data is greatly needed to further assess its distribution and viability in Montana.
Almutaster pauciflorus Alkali Marsh Aster	9/27/2017	Almutaster pauciflorus was first documented in 1988, and is now known from five sites in central and northeastern Montana. It grows in wet meadows or calcareous soil of fens within the plains.
Ligusticum verticillatum Idaho Lovage	9/27/2017	Ligusticum verticillatum occurs in northern Idaho, western Montana, and British Columbia. It has been found in Lincoln and Ravalli Counties, growing in moist forests and meadows of spruce-fir habitats, becoming common in Idaho. Herbarium specimens from Missoula and Granite Counties may be mis-identified. Current data on locations, population sizes, and threats is greatly needed.
Lobelia kalmii Kalm's Lobelia	9/27/2017	Lobelia kalmii occurs in fens and other high-organic wetlands in northwest, central, and northeast Montana. Approximately 34 observations have been made at about 23 unique locations. The central Montana location has not been observed since 1934. Current observation, population size, and threat information at documented sites is needed.
Castilleja kerryana Kerry's Paintbrush	9/27/2017	Castilleja kerryana is a recently recognized species that is found in alpine habitat within a portion of the Scapegoat Wilderness in Montana. Populations tend to be small and scattered on slopes and ridges, and apparently absent on broad, fairly flat alpine terrain. Although Castilleja species in general have brittle stems that are easily damaged by livestock, grazing is not known to occur where Kerry's Paintbrush grows. The plant appears to be limited geographically in Montana, and additional surveys are needed to accurately determine its range.
Berberis nervosa Longleaf Oregon-grape	9/27/2017	Berberis nervosa is disjunct in northern Idaho. In Montana it is known from 2-3 locations in Sanders County, of which one population in 2001 is reported to have over 1,000 plants. Additional data on locations and population sizes are greatly needed.
Triodanis leptocarpa Slim-pod Venus'-looking-glass	9/27/2017	Triodanis leptocarpa is common in the southern Great Plains and extends into eastern and central Montana. It occurs in grasslands, grass-dominated rocky slopes, and sagebrush-dominated grasslands. It has been found in grazed and ungrazed lands and appears to tolerate some disturbance. Approximately 14 locations were documented prior to 1958 and occur in central Montana. Approximately 14 locations were documented since 1974 and mostly occur in eastern Montana. Re-visits to known locations and current population data is greatly needed.
Carex glacialis Alpine Sedge	9/27/2017	Carex glacialis occurs throughout Canada, and has recently been discovered in the United States where it occurs at 4 locations in Montana. It grows in limestone fellfield habitats within the alpine. Populations are few, but appear stable. Surveys are needed to explore potential habitat, map its distribution, and determine population sizes.
Lilium columbianum Columbia Lily	9/27/2017	Lilium columbianum is currently only known from Lincoln County, where six locations have been documented in the 1980's and 1990's. This species is vulnerable to extirpation in Montana because its attractiveness, potential to be over-collected, and limited range. Native lilies have rarely survived in gardens. Current information on known locations is greatly needed.
Scolochloa festucacea Sprangletop	9/27/2017	Scolochloa festucacea occurs through most of Canada and in portions of mid-western and western States. In Montana it is known from 3 locations collected from 1949 to 1999 in Flathead County. A fourth location from a specimen with a poorly defined location in Carbon county needs to be verified. Surveys to find this species have been unsuccessful.
Lilium philadelphicum Wood Lily	9/27/2017	Lilium philadelphicum has a patchy, but wide distribution in Montana, and is often found in specialized habitats. Observations in eastern Montana have not been made since the 1930's and 1940's. This species is vulnerable to extirpation in Montana because of its attractiveness, potential to be over-collected, and habitat requirements. Native lilies have rarely survived in gardens. Current information on known locations, especially in the eastern counties, is greatly needed.
Asplenium trichomanes-ramosum Limestone Maidenhair Spleenwort	10/4/2016	Limited habitat in MT. Limited populations.
Equisetum palustre Marsh Horsetail	10/4/2016	Equisetum palustre is known from a small number of sites in seven counties of western Montana.
Equisetum pratense Meadow Horsetail	10/4/2016	Equisetum pratense has accurately been identified to occur in a few places within three counties of Montana.
Trifolium cyathiferum Cup Clover	10/4/2016	Trifolium cyathiferum occurs in two counties with limited information on population size. One occurrence was re-visited in 1998 and found to be absent due to habitat succession.
Delphinium glaucum Pale Larkspur	10/4/2016	Based on the discrepancy in the number of herbarium specimens identified as Delphinium glaucum (CPNWH 2015) and in its Montana County distribution (Lesica 2012), there seems to be an issue in how to accurately identify this species. Specimens deposited in herbaria outside of Montana will need to be examined before it can be demonstrated that this plant is more widely distributed.

ADDITIONS TO STATEWIDE LIST

SPECIES	DATE	NOTES
Delphinium depauperatum Slim Larkspur	10/4/2016	Delphinium depauperatum has been identified in Beaverhead, Flathead, and possibly Jefferson Counties in western Montana. It is found in common habitats, yet relatively few occurrences have been documented.
Trifolium microcephalum Woolly Clover	10/4/2016	Trifolium microcephalum occurs in two counties of Montana with limited population sizes.
Descurainia torulosa Wyoming Tansymustard	10/4/2016	Descurainia torulosa is known in Montana from one location in Park County; in Wyoming this species is also considered rare.
Piperia elongata Dense-flower Rein Orchid	10/4/2016	In Montana Piperia elongata is known from a single 1957 herbarium specimen collected in Lincoln County, and more recently from a few photographed specimens from Flathead, Lake, and Missoula Counties. However, the more recent observations lack data on population size and extent, habitat condition, threats, and other information. Surveys are needed to better document its status in Montana.
Allium geyeri var. geyeri Geyer's Onion	10/4/2016	In Montana this variety of Allium geyeri has been found in limited numbers with a limited distribution.
Bolboschoenus fluviatilis River Bulrush	10/4/2016	Accurate identifications of Bolboschoenus fluviatilis are found in very few populations within three counties of Montana.
Stellaria crassifolia Fleshy Stitchwort	6/18/2014	Rare in Montana where it is known from a few sparsely distributed locations.
Utricularia ochroleuca Northern Bladderwort	6/18/2014	Rare in Montana, where it is currently known from one population that may be detrimentally impacted by an adjacent gravelpit.
Senecio integerrimus var. scribneri Scribner's Ragwort	4/2/2013	Regional endemic with the core of its range in Montana. Few documented locations, though the species may be under-reported/under-collected. Some loss and degradation of habitat has likely occurred, primarily from agricultural uses.
Physaria pachyphylla Thick-leaf Bladderpod	11/5/2012	Local Endemic restricted to Carbon County and probably adjacent Big Horn County as well as adjacent WY. Currently known from only a few observations.
Pedicularis pulchella Mountain Lousewort	11/1/2012	Regional endemic from southern Montana and adjacent Wyoming with few documented locations, though the species may be under-reported/under-collected. High-elevation habitat does not appear to be at risk. Collection of additional population information may show that the viability of the species is not at risk in the state.
Mimulus clivicola North Idaho Monkeyflower	4/22/2011	Recently documented in Montana from 1 collection from 2010.
Erigeron grandiflorus Large-flower Fleabane	2/14/2011	Known in Montana from only a couple of collections.
Botrychium lunaria Common Moonwort	2/11/2011	Rare in the state. Few observation records and population levels are poorly documented.
Botrychium lanceolatum Lanceleaf Moonwort	2/11/2011	Rare in the state. Very few observation records and population levels are poorly documented.
Botrychium simplex Least Moonwort	2/11/2011	Rare in the state. Very few observation records and population levels are poorly documented.
Botrychium pinnatum Northern Moonwort	2/11/2011	Rare in the state. Very few observation records and population levels are poorly documented.
Pinus albicaulis Whitebark Pine	2/11/2011	Large declines in population levels and continued threats from white pine blister rust and mountain pine beetle attacks threaten the long-term viability of the species.
Mimulus floribundus Floriferous Monkeyflower	2/11/2011	Known in Montana from two historical collections.
Symphyotrichum molle Soft Aster	2/11/2011	Known in Montana from 1 collection from the Bighorn Mtns. Though its exact status is uncertain, its rarity warrants its inclusion as a Species of Concern.
Mimulus hymenophyllus Thinsepel monkeyflower	2/11/2011	Known in Montana from only 1 locality.
Penstemon humilis Low Beardtongue	12/16/2010	Known in Montana from 1 collection from Beaverhead County.
Douglasia conservatorium Bloom Peak Douglasia	3/16/2010	Described as a new species in 2010 based on a single location along the Idaho/Montana border.
Senecio elmeri Elmer's Ragwort	10/26/2009	Senecio elmeri is the correct identity for the single Montana location of what was previously and incorrectly called Senecio spribillei.
Physaria ludoviciana Silver Bladderpod	6/8/2009	Restricted in Montana to sandy sites in the extreme eastern portion of the state.

ADDITIONS TO STATEWIDE LIST

SPECIES	DATE	NOTES
Botrychium sp. 4 Adnate Moonwort	2/1/2008	A recently described species which is globally rare and recently discovered in northwest Montana.
Botrychium gallicomontanum Frenchman's Bluff Moonwort	2/1/2008	A recently described species which is globally rare and recently discovered in northwest Montana.
Botrychium michiganense Michigan Moonwort	2/1/2008	A recently described species which is globally rare and recently discovered in northwest Montana.
Botrychium tunux Moosewort	2/1/2008	A recently described species which is globally rare and recently discovered in northwest Montana.
Botrychium yaaxudakeit Yakutat Moonwort	2/1/2008	A recently described species which is globally rare and recently discovered in northwest Montana.
Delphinium burkei Meadow Larkspur	2/1/2008	Rare. Currently known from a few locations in western Montana in mesic meadows and grasslands.
Castilleja nivea Snow Indian Paintbrush	12/14/2007	Rare. Currently known from only a few collections from sw and south-central Montana mountain ranges. Most of these collections were made more than 30 years ago.
Cirsium pulcherrimum Wyoming Thistle	12/15/2006	
Botrychium montanum Mountain Moonwort	6/1/2006	
Collomia debilis var. camporum Alpine Collomia	6/1/2006	
Erigeron allocotus Big Horn Fleabane	6/1/2006	
Draba daviesiae Bitterroot Draba	6/1/2006	
Ipomoea leptophylla Bush morning-glory	6/1/2006	
Penstemon caryi Cary's Beardtongue	6/1/2006	
Cardamine rupicola Cliff Toothwort	6/1/2006	
Polygonum polygaloides ssp. confertiflorum Dense-flower Knotweed	6/1/2006	
Senecio eremophilus Desert Groundsel	6/1/2006	
Physaria klausii Divide Bladderpod	6/1/2006	
Erigeron flabellifolius Fan-leaved Fleabane	6/1/2006	
Castilleja crista-galli Greater Red Indian Paintbrush	6/1/2006	
Oxytropis lagopus var. conjugans Hare's-foot Locoweed	6/1/2006	
Delphinium bicolor ssp. calcicola Limestone Larkspur	6/1/2006	
Pediomelum hypogaeum var. hypogaeum Little Indian Breadroot	6/1/2006	
Camissonia subacaulis Long-leaf Evening-primrose	6/1/2006	

ADDITIONS TO STATEWIDE LIST

SPECIES	DATE	NOTES
Cirsium longistylum Long-styled Thistle	6/1/2006	
Synthyris canbyi Mission Mountain kittentails	6/1/2006	
Brickellia oblongifolia Mojave Brickellbush	6/1/2006	
Eriogonum brevicaule var. canum Parasol Buckwheat	6/1/2006	
Erigeron parryi Parry's Fleabane	6/1/2006	
Pedicularis contorta var. ctenophora Pink Coil-beaked Lousewort	6/1/2006	
Eriogonum soliceps Railroad Canyon Wild Buckwheat	6/1/2006	
Sphaeromeria capitata Rock-tansy	6/1/2006	
Physaria saximontana var. dentata Rocky Mountain Twinpod	6/1/2006	
Pedicularis crenulata Scallop-leaf Lousewort	6/1/2006	
Pedicularis contorta var. rubicunda Selway Coil-beaked Lousewort	6/1/2006	
Castilleja gracillima Slender Indian Paintbrush	6/1/2006	
Townsendia spathulata Sword Townsend-daisy	6/1/2006	
Draba crassa Thick-leaf Whitlow-grass	6/1/2006	
Penstemon flavescens Yellow Beardtongue	6/1/2006	
Calamagrostis tweedyi Cascade reedgrass	6/1/2006	
Listera borealis Northern Twayblade	6/1/2006	
Papaver pygmaeum Alpine Glacier Poppy	6/1/2001	
Salix cascadiensis Cascade Willow	6/1/2001	
Githopsis specuarioides Common Blue-cup	6/1/2001	
Physaria douglasii Douglas Bladderpod	6/1/2001	
Viola selkirkii Great-spurred Violet	6/1/2001	
Cryptantha humilis Round-headed Cryptantha	6/1/2001	
Mimulus ringens Square-stem Monkeyflower	6/1/2001	

ADDITIONS TO STATEWIDE LIST

SPECIES	DATE	NOTES
Carex chalciolepis Copper-scale Sedge	6/1/2001	Previously referred to as C. chalciolepis
Carex lacustris Lake-bank Sedge	6/1/2001	
Acorus americanus Sweetflag	6/1/2001	
Botrychium pallidum Pale Moonwort	3/1/1999	
Balsamorhiza hookeri Hooker's Balsamroot	3/1/1999	
Alnus rubra Red Alder	3/1/1999	
Erigeron tener Slender Fleabane	3/1/1999	
Mimulus ampliatus Stalk-leaved Monkeyflower	3/1/1999	Previously referred to as M. patulus
Ribes laxiflorum Trailing Black Currant	3/1/1999	
Puccinellia lemmonii Lemmon's Alkaligrass	3/1/1999	
Sisyrinchium septentrionale Northern Blue-eyed-grass	3/1/1999	
Carex pallescens Palish Sedge	3/1/1999	
Lycopodium sitchense Alaskan Clubmoss	6/1/1997	
Botrychium campestre Prairie Moonwort	6/1/1997	
Botrychium pedunculosum Stalked Moonwort	6/1/1997	
Eriogonum visherii Visher's Buckwheat	6/1/1997	
Carex chalciolepis Copper-scale Sedge	6/1/1997	Previously referred to as C. chalciolepis
Carex nelsonii Nelson's Sedge	6/1/1997	
Carex vaginata Sheathed Sedge	6/1/1997	
Evax prolifera Big-head Evax	5/1/1996	
Potentilla hyparctica Low Arctic Cinquefoil	5/1/1996	
Elatine brachysperma Short-seeded Waterwort	5/1/1996	
Eriophorum viridicarinatum Green-keeled Cottonsedge	5/1/1996	
Carex prairea Prairie Sedge	5/1/1996	
Spiranthes diluvialis Ute Ladies'-tresses	5/1/1996	

ADDITIONS TO STATEWIDE LIST

SPECIES	DATE	NOTES
Botrychium lineare Linearleaf Moonwort	5/1/1995	
Physaria brassicoides Double Bladderpod	5/1/1995	
Heterotheca villosa var. depressa Low Hairy Goldenaster	5/1/1995	
Lomatogonium rotatum Marsh Felwort	5/1/1995	
Primula incana Mealy Primrose	5/1/1995	
Lomatium nuttallii Nuttall Desert-parsley	5/1/1995	
Asclepias ovalifolia Ovalleaf Milkweed	5/1/1995	
Eustoma grandiflorum Showy Prairie-gentian	5/1/1995	
Gymnosteris parvula Small-flower Gymnosteris	5/1/1995	
Asclepias incarnata Swamp Milkweed	5/1/1995	
Poa laxa ssp. banffiana Banff Bluegrass	5/1/1995	
Trisetum orthochaetum Missoula County Oats	5/1/1995	
Scirpus pendulus Pendulous Bulrush	5/1/1995	
Poa arnowiae Short-leaved Bluegrass	5/1/1995	Previously called P. curta
Eriophorum gracile Slender Cottongrass	5/1/1995	
Botrychium ascendens Upward-lobed Moonwort	5/1/1994	
Pyrrocoma carthamoides var. subsquarrosa Beartooth Large-flowered Goldenweed	5/1/1994	
Physalis heterophylla Clammy Ground-cherry	5/1/1994	
Senecio pauciflorus Few-flowered Butterweed	5/1/1994	
Penstemon globosus Globe Beardtongue	5/1/1994	
Stellaria jamesiana James Stitchwort	5/1/1994	
Delphinium bicolor ssp. calcicola Limestone Larkspur	5/1/1994	Referrable to D. bicolor ssp. novum prior to 1995
Cryptantha humilis Round-headed Cryptantha	5/1/1994	
Townsendia leptotes Slender Townsend-daisy	5/1/1994	

ADDITIONS TO STATEWIDE LIST

SPECIES	DATE	NOTES
Ipomopsis minutiflora Small-flower Ipomopsis	5/1/1994	
Lomatium attenuatum Taper-tip Desert-parsley	5/1/1994	
Physaria didymocarpa var. lanata Woolly Twinpod	5/1/1994	
Saxifraga hirculus Yellow Marsh Saxifrage	5/1/1994	
Carex luzulina var. atropurpurea Black and Purple Sedge	5/1/1994	
Oryzopsis contracta Contracted Indian Ricegrass	5/1/1994	
Scheuchzeria palustris Pod Grass	5/1/1994	
Cyperus erythrorhizos Red-root Flatsedge	5/1/1994	
Eriophorum scheuchzeri Scheuchzer Cotton-grass	5/1/1994	
Primula alcalina Alkali Primrose	4/1/1993	
Papaver pygmaeum Alpine Glacier Poppy	4/1/1993	
Draba daviesiae Bitterroot Draba	4/1/1993	
Sphaeromeria argentea Chicken-sage	4/1/1993	
Cardamine rupicola Cliff Toothwort	4/1/1993	
Oxytropis campestris var. columbiana Columbia Locoweed	4/1/1993	
Erigeron flabellifolius Fan-leaved Fleabane	4/1/1993	
Cuscuta pentagona Field Dodder	4/1/1993	
Oxytropis lagopus var. conjugans Hare's-foot Locoweed	4/1/1993	
Cymopterus hendersonii Henderson's Wavewing	4/1/1993	
Penstemon grandiflorus Large Flowered Beardtongue	4/1/1993	
Braya humilis Low Braya	4/1/1993	
Viguiera multiflora Many-flowered Viguiera	4/1/1993	
Stenotus multicaulis Many-stem Goldenweed	4/1/1993	
Cryptantha scoparia Miner's Candle	4/1/1993	

ADDITIONS TO STATEWIDE LIST		
SPECIES	DATE	NOTES
Synthyris canbyi Mission Mountain kittentails	4/1/1993	
Nama densum Nama	4/1/1993	
Oxytropis deflexa var. foliolosa Nodding Locoweed	4/1/1993	
Eriogonum ovalifolium var. ovalifolium Oval-leaf Buckwheat	4/1/1993	Previously referred to as E. ovalifolium var. nevadense
Eriogonum brevicaule var. canum Parasol Buckwheat	4/1/1993	E. lagopus
Oxytropis parryi Parry's Locoweed	4/1/1993	
Physalis pumila ssp. hispida Prairie Ground-cherry	4/1/1993	Previously referred to as P. virginiana var. hispida
Sphaeromeria capitata Rock-tansy	4/1/1993	
Physaria saximontana var. dentata Rocky Mountain Twinpod	4/1/1993	
Draba globosa Round-fruited Draba	4/1/1993	
Claytonia arenicola Sand Springbeauty	4/1/1993	
Pedicularis contorta var. rubicunda Selway Coil-beaked Lousewort	4/1/1993	
Mimulus breviflorus Short-flowered Monkeyflower	4/1/1993	
Pediocactus simpsonii Simpson's Hedgehog Cactus	4/1/1993	
Camissonia parvula Small Camissonia	4/1/1993	
Eriogonum salsuginosum Smooth Buckwheat	4/1/1993	
Chenopodium subglabrum Smooth Goosefoot	4/1/1993	
Solidago velutina Three-nerved Goldenrod	4/1/1993	
Transberingia bursifolia ssp. virgata Twiggy Halimolobos	4/1/1993	
Symphyotrichum lanceolatum White Panicle Aster	4/1/1993	Previously referred to as Aster simplex var. ramosissimus
Polygonum polygaloides White-margin Knotweed	4/1/1993	
Penstemon flavescens Yellow Beardtongue	4/1/1993	
Muhlenbergia minutissima Annual Muhly	4/1/1993	

ADDITIONS TO STATEWIDE LIST

SPECIES	DATE	NOTES
Carex rostrata Glaucus Beaked Sedge	4/1/1993	
Phippsia algida Ice Grass	4/1/1993	
Carex eburnea Ivory Sedge	4/1/1993	
Stipa lettermanii Letterman's Needlegrass	4/1/1993	
Liparis loeselii Loesel's Twayblade	4/1/1993	
Trisetum orthochaetum Missoula County Oats	4/1/1993	
Agrostis mertensii Northern Bentgrass	4/1/1993	
Scirpus pallidus Pale Bulrush	4/1/1993	
Eriophorum callitrix Sheathed Cotton-grass	4/1/1993	
Acorus americanus Sweetflag	4/1/1993	
Juncus triglumis Three-flowered Rush	4/1/1993	
Stipa thurberiana Thurber's Needlegrass	4/1/1993	
Dichanthelium wilcoxianum Wilcox's Panic Grass	4/1/1993	

This section is not Filtered

SPECIES REMOVED FROM STATEWIDE LIST		
SPECIES	DATE	NOTES
Erigeron grandiflorus Large-flower Fleabane	9/10/2021	Erigeron grandiflorus (PDAST3M150) was combined with Erigeron simplex (PDAST3M3T0) by FNA and has a widespread distribution in relatively secure alpine habitats.
Carex multcostata Many-ribbed Sedge	9/1/2021	A statewide review of this species is warranted given changes in its nomenclature and uncertainty in the identification of many specimens.
Castilleja cervina Deer Indian Paintbrush	3/25/2021	<i>Castilleja cervina</i> is not documented in Montana (MTNHP Status Review in 2021). It was included in the <i>Flora of Montana</i> (Booth and Wright 1966) and <i>Vascular Plants of Montana</i> (Dorn 1984) possibly based on a specimen collected by R.S. Williams (1029) from "Columbia Falls, Mont." on July 10, 1894 and deposited at the Montana State University Herbarium (MONT 2775). However Mark Egger, author for the <i>Castilleja</i> treatment in the <i>Flora of North America</i> , determined that the specimen is <i>Castilleja flava</i> , which does match the identification for another Williams 1029 specimen collected on July 18, 1894 in "Columbia Falls" and deposited at the University of Montana Herbarium (MONTU 7081) (Egger pers. comm.). A specimen collected in 1990 and deposited at the Rocky Mountain Herbarium (RM 561358; Brooks 19999) is most likely <i>Castilleja flava</i> based on the location but not the specimen itself (Egger pers. comm.). <i>Castilleja cervina</i> is known from neighboring British Columbia and Alberta in Canada. If it does occur in Montana, it should be looked for in the very northwestern portion of the state (Egger pers. comm.). A conservation status rank is not applicable (SNA) because this plant is not known to occur in Montana.
Pedimelum hypogaeum var. hypogaeum Little Indian Breadroot	6/10/2013	Moved to PSOC status. Status re-determined as relatively low risk, low to moderate priority due to widespread geographic range, occurrence in over a dozen subwatersheds and low threat levels. Population numbers are small according to the limited data available, though additional surveys would likely find more populations as well as document many more individuals.
Sphaeralcea munroana White-stemmed globemallow	5/30/2013	Species was moved to PSOC status pending the collection and availability of additional information concerning the species' conservation needs and population dynamics in Montana. Most documented occurrences are from roadsides and these may be adventive or introductions.
Polygonum austiniiae Austin's Knotweed	5/29/2013	Status re-determined as relatively low risk, low to moderate priority due to widespread geographic range, occurrence in many subwatersheds, low threat levels and habitat trends that appear to be stable.
Phlox andicola Plains Phlox	5/29/2013	Status re-determined as relatively low risk, low to moderate priority due to widespread geographic range, moderate population levels, low intrinsic vulnerability and low threat levels.
Solidago velutina Three-nerved Goldenrod	5/24/2013	Species is only known in Montana from one 1980 collection in the Stillwater River Valley with little additional data available. Until additional documentation on the species distribution, abundance, habitat preferences and vulnerability becomes available, status as a Species of Concern is unwarranted.
Ranunculus hyperboreus High Northern Buttercup	5/20/2013	Status re-determined as low risk, low priority due to relatively widespread geographic range, occurrence in numerous subwatersheds and low threat levels. Additionally, the species does not appear to be restricted to rare habitats nor have intrinsic characteristics that make it especially vulnerable. See state rank details for additional information.
Sphenopholis intermedia Slender Wedgegrass	2/22/2013	Rare to uncommon in the state, where it is sporadically distributed in various mesic sites. Species may respond favorably to some disturbance and threats appear to be minimal, as such its viability in the state does appear to be at significant risk. As a result, the species was moved to the Potential Species of Concern Status pending additional information.
Balsamorhiza macrophylla Large-leaved Balsamroot	1/4/2013	Status re-determined as relatively low risk, low to moderate priority due to combination of moderate population levels, low threat levels, and habitat trends that appear to be stable. Additionally, the species does not appear to be restricted to rare habitats nor have intrinsic characteristics that make it especially vulnerable.
Botrychium montanum Mountain Moonwort	6/7/2012	Status re-determined as relatively low risk, low to moderate priority due to widespread geographic range, occurrence in many subwatersheds, low threat levels and habitat trends that appear to be stable.
Cirsium brevistylum Short-styled Thistle	6/7/2012	Dropped from SOC status pending additional information and a re-evaluation of its status to determine if the species' viability or its habitat is at risk. Unclear if the species has benefited or expanded its range from human-caused disturbances.
Botrychium lunaria Common Moonwort	6/1/2012	Status re-determined as low risk, low priority due to widespread geographic range, occurrence in numerous subwatersheds, low threat levels and habitat trends that appear to be stable. See additional state rank details.
Stellaria crassifolia Fleshy Stitchwort	5/29/2012	Species is poorly documented from Montana and its conservation priority and needs cannot be accurately assessed without additional information. Dropped from SOC status pending additional information and a re-evaluation of its status to determine if the species' viability or its habitat is at risk.
Stellaria jamesiana James Stitchwort	5/29/2012	Species is poorly documented from Montana and its conservation priority and needs cannot be accurately assessed without additional information. Dropped from SOC status pending additional information and a re-evaluation of its status to determine if the species' viability or its habitat is at risk.
Suckleya suckleyana Poison Suckleya	5/29/2012	Species is poorly documented from Montana and its conservation priority and needs cannot be accurately assessed without additional information. Dropped from SOC status pending additional information and a re-evaluation of its status to determine if the species' viability or its habitat is at risk.
Listera borealis Northern Twayblade	5/4/2012	Status re-determined as low risk, low priority due to widespread geographic range, occurrence in many subwatersheds, low threat levels and habitat trends that appear to be stable.
Juncus hallii Hall's Rush	3/12/2012	Status re-determined as low risk, low priority due to its occurrence in at least 15 subwatersheds, low threat levels, habitat trends that appear stable and overall low risk scores in all vulnerability factors.
Sphaeromeria capitata Rock-tansy	1/5/2012	Regional endemic, though population levels are robust, threats to the species' viability are minimal and large areas of intact habitat exist.
Penstemon globosus Globe Beardtongue	3/18/2011	Though rare in the state, it is more common and widespread in southwest Montana than previously reported by MTNHP. Its habitat and viability generally do not appear to be at risk in Montana.
Castilleja crista-galli Greater Red Indian Paintbrush	3/18/2011	Though uncommon in the state, it is more common and widespread in southwest Montana than previously reported by MTNHP. Its habitat and viability generally do not appear to be at risk in Montana.

SPECIES REMOVED FROM STATEWIDE LIST

SPECIES	DATE	NOTES
Potentilla uniflora One-flowered Cinquefoil	3/1/2011	Though rare in the state, the species does not appear to be at any significant risk of extirpation as a result of relatively healthy population levels and lack of threats to those populations and the species' habitat.
Poa arnowiae Short-leaved Bluegrass	3/3/2010	Moved to Status Under Review pending further taxonomic clarification of Poa anowiae in relation to Poa wheeleri and the previously used name Poa curta. Additional review of Montana material is needed.
Eustoma grandiflorum Showy Prairie-gentian	2/11/2010	Removed from SOC status due to insufficient information on the habitat and locality of the single Montana collection. May have been an isolated introduction into the state.
Townsendia spathulata Sword Townsend-daisy	9/16/2009	The species' viability in the state does not appear to be at risk due in part to its relatively widespread distribution in southwest and south-central Montana and its overall abundance.
Delphinium bicolor ssp. calcicola Limestone Larkspur	9/11/2009	A Montana endemic that is widespread in sw Montana and locally common in some habitats. The viability of this endemic subspecies does not appear to be at risk.
Orogenia linearifolia Great Basin Indian-potato	5/27/2009	More common than previously known with few potential threats to the viability of the species in MT
Ranunculus jovis Jove's Buttercup	5/27/2009	More common than previously known with very few potential threats to the viability of the species in MT
Erigeron radicans Taprooted Fleabane	4/8/2008	Removed due to overall abundance and lack of threats to high elevation habitats.
Eriogonum brevicaulis var. canum Parasol Buckwheat	12/15/2006	Locally common in parts of Carbon and Big Horn Counties.
Trifolium cyathiferum Cup Clover	6/1/2006	Status of the species in Montana requires additional review. At least 2 of the 3 documented locations in Montana are likely adventive.
Senecio pauciflorus Few-flowered Butterweed	6/1/2006	Status of the species in Montana requires additional review.
Carex chalciolepis Copper-scale Sedge	6/1/2006	Reports of this species from Montana require additional review.
Carex pallescens Palish Sedge	6/1/2006	Occurrences of this species in Montana are likely introduced.
Cypripedium parviflorum Small Yellow Lady's-slipper	6/1/2006	Moved to PSOC list due in part to the number of known occurrences, level of threat to the species and the relatively wide distribution in the state.
Cirsium longistylum Long-styled Thistle	12/15/2004	Removed from SOC status at the time as a result of review showing that a state rank of S3 was warranted.
Lycopodium sitchense Alaskan Clubmoss	4/1/2003	
Botrychium montanum Mountain Moonwort	4/1/2003	
Allotropa virgata Candystick	4/1/2003	
Chrysosplenium tetrandrum Northern Golden-carpet	4/1/2003	
Castilleja gracillima Slender Indian Paintbrush	4/1/2003	
Carex livida Pale Sedge	4/1/2003	
Senecio eremophilus Desert Groundsel	6/1/2001	S. eremophilus var eremophilus
Eurybia glauca Gray Aster	6/1/2001	
Viola renifolia Kidney-leaf White Violet	6/1/2001	

SPECIES REMOVED FROM STATEWIDE LIST

SPECIES	DATE	NOTES
Pedimelum hypogaeum var. hypogaeum Little Indian Breadroot	6/1/2001	
Salix wolfii var. wolfii Wolf Willow	6/1/2001	
Carex magellanica Poor Sedge	6/1/2001	
Botrychium minganense Mingan Island Moonwort	3/1/1999	
Salix cascadiensis Cascade Willow	3/1/1999	
Myosotis verna Early Forget-me-not	3/1/1999	
Conioselinum scopulorum Hemlock Parsley	3/1/1999	
Helenium hoopesii Orange Sneezeweed	3/1/1999	
Cryptantha flavoculata Pale Yellow Cryptantha	3/1/1999	
Agoseris lackschewitzii Pink Agoseris	3/1/1999	
Gentiana prostrata Pygmy Gentian	3/1/1999	
Cryptantha humilis Round-headed Cryptantha	3/1/1999	
Gentianella tenella Slender Gentian	3/1/1999	
Halenia deflexa Spurred Gentian	3/1/1999	
Bidens comosa Three-lobed Beggarticks	3/1/1999	
Carex neurophora Alpine Nerved Sedge	3/1/1999	
Calamagrostis tweedyi Cascade reedgrass	3/1/1999	
Carex chalciolepis Copper-scale Sedge	3/1/1999	Previously referred to as C. chalciolepis
Allium fibrillum Fringed Onion	3/1/1999	
Carex nelsonii Nelson's Sedge	3/1/1999	
Agrostis mertensii Northern Bentgrass	3/1/1999	
Juncus triglumis Three-flowered Rush	3/1/1999	
Papaver pygmaeum Alpine Glacier Poppy	6/1/1997	
Evax prolifera Big-head Evax	6/1/1997	
Physaria klausii Divide Bladderpod	6/1/1997	

SPECIES REMOVED FROM STATEWIDE LIST

SPECIES	DATE	NOTES
Erigeron flabellifolius Fan-leaved Fleabane	6/1/1997	
Cuscuta pentagona Field Dodder	6/1/1997	
Heterotheca villosa var. depressa Low Hairy Goldenaster	6/1/1997	Chrysopsis villosa
Eriogonum brevicaulis var. canum Parasol Buckwheat	6/1/1997	E. lagopus
Spiraea x pyramidata Pyramidal Spiraea	6/1/1997	
Erigeron flagellaris Running Fleabane	6/1/1997	
Pedicularis contorta var. rubicunda Selway Coil-beaked Lousewort	6/1/1997	
Madia minima Small-headed Tarweed	6/1/1997	
Bidens vulgata Tall Bur-marigold	6/1/1997	Specifically B. vulgata var. schizantha
Symphyotrichum lanceolatum White Panicle Aster	6/1/1997	Previously referred to as Aster simplex var. ramosissimus
Polygonum polygaloides White-margin Knotweed	6/1/1997	
Lilium columbianum Columbia Lily	6/1/1997	
Oryzopsis contracta Contracted Indian Ricegrass	6/1/1997	
Eriophorum viridicarinatum Green-keeled Cottonsedge	6/1/1997	
Carex eburnea Ivory Sedge	6/1/1997	
Trisetum orthochaetum Missoula County Oats	6/1/1997	
Scirpus pendulus Pendulous Bulrush	6/1/1997	
Astragalus platytropis Broad-keeled Milkvetch	5/1/1996	
Penstemon caryi Cary's Beardtongue	5/1/1996	
Castilleja pilosa var. longispica Parrot-head Indian Paintbrush	5/1/1996	C. longispica
Physalis pumila ssp. hispida Prairie Ground-cherry	5/1/1996	Previously referred to as P. virginiana var. hispida
Carex luzulina var. atropurpurea Black and Purple Sedge	5/1/1996	
Carex torreyi Torrey's Sedge	5/1/1996	
Erigeron allocotus Big Horn Fleabane	5/1/1995	Regional endemic, secure

SPECIES REMOVED FROM STATEWIDE LIST		
SPECIES	DATE	NOTES
Draba daviesiae Bitterroot Draba	5/1/1995	Regional endemic, secure
Physalis heterophylla Clammy Ground-cherry	5/1/1995	Adventive
Cardamine rupicola Cliff Toothwort	5/1/1995	State endemic, secure
Astragalus chamaeleuce Ground Milkvetch	5/1/1995	Many populations, low threats
Oxytropis lagopus var. conjugans Hare's-foot Locoweed	5/1/1995	State endemic, secure
Cymopterus hendersonii Henderson's Wavewing	5/1/1995	Taxonomic revision pending
Delphinium bicolor ssp. calcicola Limestone Larkspur	5/1/1995	Referable to D. bicolor ssp. novum prior to 1995
Ericameria discoidea var. linearis Linear-leaved Whitestem Goldenbush	5/1/1995	Many populations, low threats
Stenotus multicaulis Many-stem Goldenweed	5/1/1995	New populations, low threats
Synthyris canbyi Mission Mountain Kittentails	5/1/1995	Regional endemic, secure
Sphaeromeria capitata Rock-tansy	5/1/1995	Many populations, low threats
Physaria saximontana var. dentata Rocky Mountain Twinpod	5/1/1995	
Epilobium suffruticosum Shrubby Willowherb	5/1/1995	Many populations, low threats
Gaultheria ovatifolia Slender Wintergreen	5/1/1995	Many populations, low threats
Lorandersonia linifolia Spearleaf Rabbitbrush	5/1/1995	Locally common, low threats
Townsendia spathulata Sword Townsend-daisy	5/1/1995	Many populations, low threats
Trifolium latifolium Twin Clover	5/1/1995	Many populations, low threats
Trifolium microcephalum Woolly Clover	5/1/1995	Many populations, low threats
Penstemon flavescens Yellow Beardtongue	5/1/1995	Regional endemic, secure
Muhlenbergia minutissima Annual Muhly	5/1/1995	Many populations, low threats
Eriophorum viridicarinum Green-keeled Cottonsedge	5/1/1995	Many populations, locally common
Amphiscirpus nevadensis Nevada Clubrush	5/1/1995	Many populations, low threats
Scirpus pallidus Pale Bulrush	5/1/1995	Many populations, low threats
Dichanthelium acuminatum Panic Grass	5/1/1995	Many populations, low threats. Previously referred to as Panicum occidentale

SPECIES REMOVED FROM STATEWIDE LIST

SPECIES	DATE	NOTES
Acorus americanus Sweetflag	5/1/1995	Specimen review needed
Stipa thurberiana Thurber's Needlegrass	5/1/1995	Probably accidental
Carex vallicola Valley Sedge	5/1/1995	Many populations, low threats
Dichanthelium wilcoxianum Wilcox's Panic Grass	5/1/1995	Many populations, low threats
Lycopodium alpinum Alpine Clubmoss	5/1/1994	More common than previously known
Orobanche corymbosa Flat-topped Broomrape	5/1/1994	More common than previously known
Stanleya viridiflora Green Prince's plume	5/1/1994	Limited distribution
Arenaria kingii King's Arenaria	5/1/1994	More common than previously known
Eriogonum ovalifolium var. ovalifolium Oval-leaf Buckwheat	5/1/1994	More common than previously known. Previously referred to as E. ovalifolium var. nevadense
Astragalus leptaleus Park Milkvetch	5/1/1994	Limited distribution
Castilleja flava var. rustica Rustic Indian Paintbrush	5/1/1994	More common than previously known. Many populations, low threats
Astragalus argophyllus Silver-leaved Milkvetch	5/1/1994	More common than previously known
Pediocactus simpsonii Simpson's Hedgehog Cactus	5/1/1994	More common than previously known
Erigeron gracilis Slender Fleabane	5/1/1994	More common than previously known
Mimulus suksdorfii Suksdorf Monkeyflower	5/1/1994	More common than previously known
Senecio debilis Weak Groundsel	5/1/1994	Limited distribution
Trisetum orthochaetum Missoula County Oats	5/1/1994	Sterile hybrid
Selaginella watsonii Watson's Spikemoss	4/1/1993	More common than previously known
Ipomopsis pumila Dwarf Ipomopsis	4/1/1993	More common than previously known
Ligusticum filicinum Fern-leaf Lovage	4/1/1993	More common than previously known
Gilia leptomeria Great Basin Gilia	4/1/1993	More common than previously known
Townsendia incana Hoary Townsend-daisy	4/1/1993	More common than previously known
Geocaulon lividum Northern Toadflax	4/1/1993	More common than previously known
Claytonia multiscapa Rydberg's Springbeauty	4/1/1993	1994 note: More common than previously known
Camissonia minor Small-flowered Evening-primrose	4/1/1993	More common than previously known

SPECIES REMOVED FROM STATEWIDE LIST

SPECIES	DATE	NOTES
Phacelia ivesiana var. glandulifera Sticky Scorpion-weed	4/1/1993	More common than previously known
Streptanthella longirostris Streptanthella	4/1/1993	More common than previously known
Gilia tweedyi Tweedy's Gilia	4/1/1993	More common than previously known. Previously referred to as G. inconspicua var. tweedyi
Xylorhiza glabriuscula Woody Aster	4/1/1993	More common than previously known
Stanleya tomentosa Woolly Prince's plume	4/1/1993	More common than previously known
Scirpus cyperinus Woolgrass	4/1/1993	Adventive

Citation for data on this website:

Montana Plant Species of Concern Report. Montana Natural Heritage Program. Retrieved on 12/7/2021, from mtnhp.org/SpeciesOfConcern/?AorP=p